## Math 1375/6578, Fall 2012

## Midterm Exam #1 - March 4, 2012

- For full credit you must show all your work.
- Begin each problem on a separate sheet of paper.
- You are allowed to use a graphing calculator or app. (But note that the exam can be completed without one).
- Use the provided sheet of coordinate planes for your graphs.
- Write your name at the top of every sheet of paper.
- 1. Consider the function f(x) = |2x 3|.
  - (a) Solve the equation |2x 3| = 5. (Extra credit: Check your answer by plugging your solution(s) back into f(x).)
  - (b) Sketch the graph of f(x).
  - (c) Label the point(s) on the graph corresponding to the solution(s) you found for the equation in part (a).
- 2. Consider the function g(x) = mx + b where m and b are constants.
  - (a) Describe in a sentence or two what the graph of g(x) looks like. In particular, describe what the quantities m and b represent on the graph.
  - (b) Sketch the graph of g(x) for  $m = -\frac{1}{2}$  and b = -1. Label the x- and y- intercepts.
- 3. Consider the piecewise-defined function h(x):

$$h(x) = \begin{cases} x^2 & \text{if } -3 < x \le 0\\ 1/x & \text{if } x > 0 \end{cases}$$

- (a) Find each of the following:
  - i. h(-2)
  - ii. h(0)
  - iii. h(100)
- (b) Sketch the graph of h(x).
- (c) What are the domain and range of h? Use interval notation or inequalities to represent these sets of real numbers.
- 4. Consider the function f(t) = 3t + 1.
  - (a) Find f(t+h) (i.e., write down the rule for computing f(t+h). Hint: Your answer will be in terms of t and h.)
  - (b) Show that the difference quotient for f simplifies as follows:

$$\frac{f(t+h) - f(t)}{h} = 3$$

(c) (Extra credit) For g(x) = mx + b (as in Exercise (2) above), show that:

$$\frac{g(x+h) - g(x)}{h} = m$$