Take Charge

of Your

DIABETES





U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES

Centers for Disease Control and Prevention National Center for Chronic Disease Prevention and Health Promotion



This book belongs to
If found, please call or write me at

Take Charge of Your Diabetes

4th Edition

— 2007 —

DEPARTMENT OF HEALTH AND HUMAN SERVICES

Centers for Disease Control and Prevention

National Center for Chronic Disease Prevention and Health Promotion

Division of Diabetes Translation

For more information

The Centers for Disease Control and Prevention's (CDC's) Diabetes Program supports diabetes control programs in all states, the District of Columbia, and eight U.S. territories or island jurisdictions. You may contact your local program for more information on diabetes.

To learn more, call toll free 1-800-CDC-INFO (232-4636), E-mail cdcinfo@cdc.gov. Information available in English, and Spanish 24 Hours/Day, 7 Days/Week.

This book is in the public domain. Anyone may reproduce any or all of the contents. It is also available on the Internet at www.cdc.gov/diabetes, or you can call toll free 1-800-CDC-INFO (232-4636) for more information or a copy of this book and others. This book is also available in Spanish.


Links or references to nonfederal organizations mentioned in this book or in the resource list are provided solely as a service to our users. These links and references do not constitute an endorsement of these organizations or their programs by CDC or the federal government, and none should be inferred. CDC is not responsible for the content of the individual organization Web pages found at these links.

Suggested citation:

Centers for Disease Control and Prevention. *Take Charge of Your Diabetes*. 4th edition. Atlanta:

U.S. Department of Health and Human Services, 2007.

This book is dedicated to all people living with diabetes, in honor of your struggles and your strength.



Many of us strive for a sense of balance in our lives. We want to keep our goals in harmony with our minds and souls. People with diabetes often have stories to share about their struggles for balance and harmony in their lives. We can honor people by listening to and learning from their stories to find meaning and hope for our own lives.

An old, well-loved story, told around the world, is about the turtle and a sure-footed animal, like a rabbit. In this story, the turtle tricks the other animal to win a race—simply by not giving up and by staying on its path. And it has to stick its neck out! It takes determination like that to face diabetes, day after day, reminding yourself that you can do it if you stick to it!

There are about 250 kinds of turtles, and almost all have the same pattern on their top shell—13 plates that fit together in harmony and balance to form a strong shell. The turtle and its shell can remind us of the harmony and balance we seek in all parts of our lives—including living with diabetes.

Contents

Sor	ne Words of Thanks vii
1	Introduction
2	Controlling Your Diabetes
3	Keeping Track of Your Blood Glucose 21
4	Feelings About Having Diabetes 41
5	Eye Problems
6	Kidney Problems 47
7	Heart and Blood Vessel Problems 51
8	Nerve Damage
9	Foot Problems
10	Dental Disease 65
11	Vaccinations
12	Pregnancy and Women's Health 73
RE	CORDS
	Records for Sick Days 79
	Tests and Goals for Each Visit 89
	Tests and Goals for Each Year 99
	Glucose Log Sheets
	Your Health Care Team 109
GL	OSSARY
RE	SOURCES 127

Some Words of Thanks

This guide was written by staff in the Centers for Disease Control and Prevention's (CDC's) Division of Diabetes Translation. The division is part of the National Center for Chronic Disease Prevention and Health Promotion, Department of Health and Human Services. We work with partners who share our mission to reduce the burden of diabetes in communities.

William H. Herman, MD, MPH, was the general editor of the first book, *Take Charge of Your Diabetes: A Guide for Care*, printed in 1991. We asked people with diabetes who read the first book to help us make the second book even more useful. For this fourth edition of the book, we have updated the scientific facts, resources, and art.


The American Association of Diabetes Educators did a survey among people with diabetes and diabetes educators to learn what people liked and didn't like about the first book. Focus groups made up of people with diabetes were held by the Health Promotion Council of SE Pennsylvania and Casals and Associates of Washington, DC. The groups gave us valuable input to help us make later books more useful.

Important support for this book's emphasis on glucose control came from the Diabetes Control and Complications Trial. Conducted by the National Institute of Diabetes and Digestive and Kidney Diseases, National Institutes of Health, this important study provided scientific proof that glucose control can help prevent or delay complications of diabetes.

Dawn Satterfield, PhD, RN, CDE and Patricia Mitchell of the CDC diabetes division's Health Communications Section were the lead writers of this book. Ann Constance, Claudia Martinez, Hope Woodward, Margaret Fowke, Nancy Haynie-Mooney, Melinda Salmon, Mike Engelgau, and the Chattahoochee Nature Center also helped with the writing. Rick Hull, Diana Toomer, Melissa Stankus, and Kristina Ernst reviewed and edited the final version of this guide. Most of the drawings were provided by the Public Health Practice Program Office, CDC, some of which were modified for the book Take Charge of Your Diabetes: A Shortened Overview for Pacific Basin Island Populations through the National Diabetes Education Program. Cygnus Corporation assisted with design and layout of the second edition. Further publication support for the third edition was provided by Palladian Partners, Inc., under Contract 200-98-0415 for the National Center for Chronic Disease Prevention and Health Promotion, CDC, Department of Health and Human Services.

1 Introduction

Diabetes touches almost every part of your life. It's a serious, lifelong condition, but there's a lot you can do to protect your health. You can take charge of your health—not only for today, but for the coming years.



Take charge of your health—for today and for the coming years.

Diabetes can cause

health problems over time. It can hurt your eyes, your kidneys, and your nerves. It can lead to problems with the blood flow in your body. Even your teeth and gums can be harmed. Diabetes in pregnancy can cause special problems. Many of these problems don't have to happen. You can do a lot to prevent them, and there are people in your community who can help. This book can help you find how to get the help you need to prevent problems.

Today and every day, strive to balance your food, physical activity, and medicine. Test your own blood glucose (also called blood sugar) to see how this balance is working out. Then make choices that help you feel well every day to protect your health.

Feeling healthy can allow you to play a big part in the life of your family and community. You may even want to join a community group in which people share their stories and help others deal with their diabetes.



A community group can help make life better for people living with diabetes.

Take Charge of Your Diabetes was written to help you take important steps to prevent problems caused by diabetes. You'll learn many useful things:

- What problems diabetes can cause.
- How to work with a health care team to prevent problems.
- Why it is important to get your blood glucose and blood pressure closer to normal.
- How to find out about resources in your community to help you prevent problems.

It's important to work with a primary health care provider, as well as other members of a team who




Work with your health care team to take charge of your diabetes.

care about your health. To find out about resources in your community, contact one of the groups listed below:

- Diabetes organizations, listed on pages 127–129 of this book.
- Local diabetes programs or hospitals.
- Your state health department's diabetes prevention and control program, which you can find by calling 800-CDC-INFO.

Ask your health care team to look over this book with you. Stay in touch with them so you will know the latest news about diabetes care.

Balance is the key word in living well with diabetes. Strive for balance in all parts of your life. With the support of your family and friends, your health care team, and your community, you can take charge of your diabetes.



Who Is This Book For?

This book was mainly written for people who found out they had diabetes as an adult. You should use it along with other information your health care providers give you.

If you've just learned you have diabetes, you'll need more details than you'll find in this book. Ask your health care provider for help. See the list beginning on page 127 for phone numbers, addresses, and web sites of organizations where you can get more information. Find out as much as you can about the three most important things for controlling your diabetes: food, physical activity, and diabetes medicine.



Your health care providers can tell you more about the topics in this book.

How to Use This Book

When you're reading this book, note these points:

- Words in **bold print** are explained in the glossary, which starts on page 117.
- The forms at the back of this book can help you and your health care team keep records of your care.
- On pages 127–129, you'll find a list of health organizations that you can call, write, or E-mail for more information about diabetes.
- When we say "health care team," we include all the people who work with you to help manage your diabetes: primary doctor, dietitian, nurse, diabetes educator, counselor, foot doctor, eye doctor, dentist, pharmacist, community health worker, and others.
- The chapters in this book deal with many topics. You may first want to read the parts that deal with your own special concerns. Take your time reading this book. There's a lot to read, but you don't have to read it all at once.

Keeping Records

You can use this book to keep some records about your health. The forms to write down details about your health begin on page 79. You can cut out these pages to take with you on your diabetes care visits. You may also want to make extra copies to use in the future. Go over these records often with your health care team. Keeping track of your health is one of the ways you can work together to control your diabetes.

On page 109, write down the names and telephone numbers of your health care team. There's enough room on these pages to write questions and other points you want to remember when you go to your visits every 4 to 6 months. On page 116, you may want to write down some contacts for community groups that deal with diabetes.

Your Health Care Providers
Primary Doctor
Name: <i>Dr. B. Harper</i>
Telephone number: 222-222-2222
Your questions:
What was my last A1C result?
When is my next eye exam due?
Important points:
— Aim for glacose 90-130 before eating,
—Less 180 1-2 hours after beginning to eat.
— Check my feet every day,

Write down the things you want to discuss with your health care team.

What Is Diabetes?

Most of the food we eat is turned into **glucose** (sugar) for our bodies to use for energy. The **pancreas**, an organ near the stomach, makes a **hormone** called **insulin** to help glucose get into our body cells. When you have **diabetes**, your body either doesn't make enough insulin or can't use its own insulin very well. This problem causes glucose to build up in your blood.

Diabetes means that a person's blood sugar is too high. Your blood always has some sugar in it because the body needs sugar for energy to keep you going. But too much sugar in the blood can cause serious damage to the eyes, kidneys, nerves, and heart.

Signs and Symptoms of Diabetes

You may recall having some of these signs before you found out you had diabetes:

- Being very thirsty.
- Urinating a lot—often at night.
- Having blurry vision from time to time.
- Feeling very tired much of the time.
- Losing weight without trying.
- Having very dry skin.
- Having sores that are slow to heal.

- Getting more infections than usual.
- Losing feeling or getting a tingling feeling in the feet.
- Vomiting.

Types of Diabetes

There are two main types of diabetes:

- **■ Type 1**.
- **■** Type 2.

Another type of diabetes appears during pregnancy in some women. It's called **gestational diabetes**. See page 75 to learn more about this type of diabetes.

One out of 10 people with diabetes has type 1 diabetes. These people usually find out they have diabetes when they are children or young adults. People with type 1 diabetes must **inject insulin** every day to live. The pancreas of a person with type 1 makes little or no insulin. Scientists are learning more about what causes the body to attack its own **beta cells** of the pancreas (an **autoimmune process**) and stop making insulin in people with certain sets of genes.



Whether you have type 1 or type 2 diabetes, work closely with your health care provider.

Most people with diabetes—9 out of 10—have type 2 diabetes. The pancreas of people with type 2 diabetes keeps making insulin for some time, but the body can't use it very well. Most people with type 2 find out about their diabetes after age 30 or 40.

Certain **risk factors** make people more likely to develop type 2 diabetes. Some of these are

- A family history of diabetes.
- Lack of exercise.
- Weighing too much.
- Being of African American, American Indian, Alaska Native, Hispanic/Latino, or Asian/Pacific Islander heritage.
- Gestational diabetes history.

You can help manage your diabetes by controlling your weight, making healthy food choices, and getting regular physical activity. Ask for help from your health care team. Some people with type 2 diabetes may also need to take **diabetes pills** or **insulin** shots to help control their diabetes.

Some people with diabetes are concerned about their family members getting diabetes. A national study show's that people may be able to prevent or delay the onset of type 2 diabetes. To find out more, talk to your health care provider, visit the CDC Diabetes Web site at www.cdc.gov/diabetes, or call 1-800-CDC-INFO.



Whether you have type 1 or type 2 diabetes, learn what your community has to offer you.

2 Controlling Your Diabetes

There's good news for people with diabetes. Studies show that keeping your **blood glucose** (also called **blood sugar**) close to normal helps prevent or delay some diabetes problems.

Through careful control, many problems such as eye disease, kidney disease, heart disease, nerve damage, and serious foot problems can be prevented or slowed. People who have **type 1 diabetes** as well as people who have **type 2 diabetes** can benefit by keeping their blood glucose levels closer to normal.

You can learn more about diabetes and ways to help you control your blood sugar by calling the National Diabetes Education Program (NDEP) at 1-800-438-5383.



You may find that your community supports your efforts to control your diabetes.

Keeping a Balance



As the turtle makes steady progress, so too must those with diabetes continue to maintain healthy lifestyles and stick to daily routines that involve regular exercise, good nutrition, glucose monitoring, and regular visits to health care providers.

To keep your glucose at a healthy level, you need to keep a balance between three important things:

- What you eat and drink.
- How much physical activity you do.
- What diabetes medicine you take (if your doctor has prescribed **diabetes pills** or **insulin**).

This book gives you only some of the facts you need. Your health care team can give you more.

A Few Things About Food

Here are some tips for making healthy eating choices:


■ *Eat regular meals*. Ask your health care team to help you choose a **meal plan**. Your dietitian may suggest you eat three meals and a snack or two every day at about the same times. Eating every 4 to 5 hours can help control blood sugar.

- *Eat a variety of foods*. Choose a variety of foods to eat so that your body gets the nutrition it
- *Eat less fat.* Avoid fried foods. Foods that are baked, broiled, grilled, boiled, or steamed are more healthy to eat. Eat meats that have little fat. When you eat dairy products (cheese, milk, yogurt, and others), choose those that have little or no fat or cream.
- *Eat less sugar*. You may find that eating less sugar helps you control your blood glucose level. Here are some things you can do to eat less sugar:
 - Eat more high-fiber foods, like vegetables, dried beans, fruit, and whole grain breads and cereals.



Choose to eat a variety of healthy foods, such as fruit and vegetables.

Ask your market to carry more heart-healthy foods.



- Drink water and other drinks that have no added sugar.
- Eat fewer foods that have extra sugar, such as cookies, cakes, pastries, candy, brownies, and sugared breakfast cereals.
- Talk with your health care team about ways to sweeten food and drinks without using sugar.

See pages 28–32 for more on ways to prevent problems when your blood glucose levels are too high or too low.

- *Eat less salt*. Eating less salt may help control your blood pressure. Here are some ways to eat less salt:
 - Use less salt when you prepare foods.
 - Cut down on processed foods, such as foods

you buy in cans and jars, pickled foods, lunch meats ("cold cuts"), and snack foods, such as chips.

- Taste your food first before adding salt. You may not need to add any.
- Use herbs and spices instead of salt to flavor your food.
- A word about drinking alcohol: Alcohol can cause health problems, especially for people with diabetes. It adds calories and doesn't give your body any nutrition. Drinking alcohol may cause dangerous reactions with medicines you take. Your blood glucose can go down too low if you drink beer, wine, or liquor on an empty stomach. If you want to include a drink in your food plan once in a while, ask your health care team how to do so safely.

A Few Things About Physical Activity

■ *It's important to be active*. Physical activity has many benefits. It can help you control your blood glucose and your weight. Physical activity

can help prevent heart and blood flow problems. Many people say they feel better when they get regular exercise.


- Start with a little. If you haven't been doing any physical activity, talk to your health care team before you begin. Walking, working in the yard, and dancing are good ways to start. As you become stronger, you can add a few extra minutes to your physical activity. If you feel pain, slow down or stop and wait until it goes away. If the pain comes back, talk with your health care team right away.
- Do some physical activity every day. It's better to walk 10 or 20 minutes each day than one hour once a week.
- Choose an activity you enjoy.

 Do an activity you really like.

 The more fun it is, the more likely you will do it each day.

 It's also good to exercise with a family member or friend.

Brisk walk is an activity almost everyone can do.



If you're already active now, but want to become more active, talk to your health care team about a safe exercise plan.

A Few Things About Diabetes Medicine

If you take diabetes pills or insulin injections to control your diabetes, ask your health care provider to explain how these work. It's important to know how and when to take diabetes medicine. If you take other medicines that are sold with or without a prescription, ask your doctor how these can affect your diabetes control. When you take insulin injections or diabetes pills, your blood glucose levels can get too low. See pages 28–32 for how to prevent levels that are too low or too high.

If you inject insulin, your health care team should be able to tell you:

- How to give yourself injections.
- When you need to change your insulin dose.

Be sure you know how and when to take your diabetes medicine.



3 Keeping Track of Your Blood Glucose

It's important to your health to control your **blood glucose** (also called **blood sugar**). Keeping your glucose level close to normal helps prevent or delay some diabetes problems, such as eye disease, kidney



Keep a daily record of your blood glucose levels.


disease, and nerve damage. One thing that can help you control your glucose level is to keep track of it. You can do this by:

- Checking your own glucose a number of times each day (**self-monitoring blood glucose**). Many people with diabetes check their glucose 2 to 4 times a day.
- Getting an **A1C** test from your health care provider about every 3 months.

You'll learn more about these tests on the next pages. These tests can help you and the rest of your diabetes health care team—doctor, diabetes educator, and others—work together to help you control your blood glucose.

Checking Your Blood Glucose Each Day

You can do a test to find out what your blood glucose is at any moment. Your health care team can show you how to do the test yourself. Using a finger prick, you place a drop of blood on a special coated strip, which "reads" your blood glucose. Many people use an electronic meter to get this reading.



Checking your own blood glucose levels is a key to taking charge of your diabetes.

Blood glucose testing can help you understand how food, physical activity, and diabetes medicine affect your glucose levels. Testing can help you make day-to-day choices about how to balance these things. It can also tell you when your glucose is too low or too high so that you can treat these problems.

Ask your health care team to help you set a goal for your glucose range and show you how to record your glucose readings in a logbook or record sheet. If you need a daily logbook, ask your health care provider for one. Or you can make copies of page

108 if you take insulin or page 106 if you don't take insulin. A sample log sheet is filled out to show you how to use each.

Be sure to write down each glucose reading and the date and time you took it. When you review your records, you can see a pattern of your recent glucose control. Keeping track of your glucose on a day-to-day basis is one of the best ways you can take charge of your diabetes.

D	Daily Log			SAMPLE				W	eek	Starti	ng <u>May 23, 2001</u>
	Bre	akfast	L	ınch	Di	nner	Bedtime Other		her		
	Dose	Blood	Dose	Blood Sugar	esog	Blood Sugar	веод	Blood Sugar	Dose	Blood Sugar	Notes
Mon		108		118		124		112			
Tues		112		109				¥- 151			* Missed evening walk. Start back tomorrow!
Wed		125		122		130		121			
Thurs		114		129		185		242			FSIckw/flu? Drinking diet soda Ketones negative.
Fri		156		148		135		130			Feeling better today
Sat		128				125		* 151		129 /1 pm	#Extrajuice made sugargo UP.
Sun		120		119		168		133			V Lunch at church

Think of your daily log sheet as a diary for taking charge of your diabetes.

Getting a Summary Lab Test (A1C)

By performing an A1C test, health provides can sum up your diabetes control for the past few months. An A1C test measures how much glucose has been sticking to your red blood cells. Since each red blood cell is replaced by a new one every 3 to 4 months, this test tells you how high the glucose levels have been during the life of the cells.

If most of your recent blood glucose readings have been near normal (70 to 140 milligrams per deciliter or mg/dL, with the higher reading mainly after meals), the A1C test will be near normal (usually about 6%–7%). If you've had many readings above normal, the extra glucose sticking to your red blood cells will make your A1C test read higher.

You should get an A1C test every 3 months if your test results are not yet at goal. You should get an AIC test at least 2 times a year if your AIC results are at goal. Ask your health care provider for the results and record them on page 91.

Ask your team to tell you the normal range of values and help you set a goal for yourself. Write your goal down on page 91 of this guide.

If your A1C test results are high, work with your team to adjust your balance of food, physical activity, and diabetes medicine. When your A1C test result is near your goal, you'll know you've balanced things well.

Have you	r health care (Record da				s with you.	
Tests and Goals			Dates ar	nd Results		
Blood Glu <u>cose Imgfell</u>	2/1/2900	6/11/2000	9/28/2000	1/5/2001	4/3/2001	
A1 c Test/Goal (%)	9.0	89	8.9	nesa /	8.2	
Weight/Goal (pounds)	180	115	112	100	165	
Blood Pressure goal:/(mm Hg)	140/90	110/86	138/20	136/82	121/80	
Foot Check	1	1	1	1	1	

Use your A1C test to track your glucose control.

Having Problems with Low Blood Glucose

In general, a blood glucose reading lower than 70 mg/dL is too low. If you take insulin or **diabetes pills**, you can have **low blood glucose** (also called **hypoglycemia**). Low blood glucose is usually caused by eating less or later than usual, being more active than usual, or taking too much diabetes medicine. Drinking beer, wine, or liquor may also cause low blood glucose or make it worse.

Low blood glucose happens more often when you're trying to keep your glucose level near normal. This is no reason to stop trying to control your diabetes. It just means you have to watch more carefully for low levels. Talk this over with your health care team.

Signs of Low Blood Glucose

Some possible signs of low blood glucose are feeling nervous, shaky, or sweaty. Sometimes people just feel tired.

The signs may be mild at first. But a low glucose level can quickly drop much lower if you don't treat it. When your glucose level is very low, you may get confused, pass out, or have seizures.

If you have any signs that your glucose may be low, test it right away. If it's less than 60 to 70 mg/dL, you need to treat it right away. See below for ways to treat low blood glucose.



If you have signs that your blood glucose is low but you can't test right then...



go ahead and treat it! (See page 26.)



Treat it every 15 minutes, until your glucose level is normal.

Treating Low Blood Glucose

If you feel like your blood glucose is getting too low but you can't test it right then, play it safe—go ahead and treat it. Eat 10 to 15 grams of **carbohydrate** right away. See the box below for examples of foods and liquids with this amount of carbohydrate.

Foods and Liquids for Low Blood Glucose (each item has 15 grams of carbohydrate)					
Food Item Amount					
Sugar packets	3 to 4				
Fruit juice	¹ / ₂ cup (4 ounces)				
Soda pop (not diet)	¹ / ₂ cup (4 ounces)				
Hard candy	3 to 5 pieces				
Sugar or honey	4 teaspoons				
Glucose tablets 3 to 4					

Check your blood glucose again in 15 minutes. Eat another 10 to 15 grams of carbohydrate every 15 minutes until your blood glucose is above 70 mg/dL.

Eating or drinking an item from the list on this page will keep your glucose up for only about 30 minutes. So if your next planned meal or snack

is more than 30 minutes away, you should go ahead and eat a small snack, something like crackers and a tablespoon of peanut butter.

In your glucose logbook or record sheet, write down the numbers and the times when low levels happen. Think about what may be causing them. If you think you know the reason, write it beside the numbers you recorded. You may need to call your health care provider to talk about changing

medicine.

Tell family members, close friends, teachers, and people at work that you have diabetes. Tell them how to know when your blood glucose is low. Show them what to do if you can't treat yourself. Someone will need to give you fruit juice, soda pop (not diet), or sugar.

your diet, activity, or diabetes

Write in your logbook why you had a low blood glucose level.

If you can't swallow, someone will need to give you a shot of **glucagon** and call for help. Glucagon is a prescription medicine that raises the blood glucose and is injected like insulin. If you take insulin, you should have a glucagon kit handy. Teach family members, roommates, and friends when and how to use it.

Waiting to treat low blood glucose is not safe. You may be in danger of passing out. If you get confused, pass out, or have a seizure, you need emergency help. Don't try to drive yourself to get help. Be prepared for an emergency.



In a low blood glucose emergency, you may need to go to the hospital.

Preventing Low Blood Glucose

Keep a balance

Try to stay close to your usual schedule of eating, activity, and medicine. If you're late getting a meal or if you're more active than usual, you may need an extra snack. See page 37 for more ideas about managing your diabetes.

Check your blood glucose

Keeping track of your blood glucose is a good way to know when it tends to run low. Show your logbook or record sheet to your health care providers. Be sure to let them know if you're having a number of low blood glucose readings a week. To be safe, always check your blood glucose before doing any of these things:

- Driving a vehicle.
- Using heavy equipment.
- Being very physically active.
- Being active for a long time.

Ask your health care team whether you should check your glucose before (or during) any other activities. Write these in the space below.

Be prepared


Always carry some type of carbohydrate sugar food or drink with you so you'll be ready at any time to treat a low glucose level. See the list on page 26 for snacks that have 15 grams of carbohydrate.



Always carry along some food with carbohydrates in it.

Always wear something (like an identification bracelet) that says you have diabetes. Carry a card in your wallet that says you have diabetes and tells if you use medicine to treat it.

Wear something that lets others know you have diabetes, in case of an emergency.



Having Problems with High Blood Glucose

For most people, blood glucose levels that stay higher than 140 mg/dL (before meals) are too high. Talk with your health care team about the glucose range that is best for you.

Eating too much food, being less active than usual, or taking too little diabetes medicine are some common reasons for **high blood glucose** (or **hyperglycemia**). Your blood glucose can also go up when you're sick or under stress.

Over time, high blood glucose can damage body organs. For this reason, many people with diabetes try to keep their blood glucose in balance as much as they can.

Some people with type 2 diabetes may not feel the signs of high blood glucose until their blood glucose is higher than 300. People with blood glucose higher than 300 are more likely to have **dehydration**. Dehydration can become a serious problem if not treated right away.

Your blood glucose is more likely to go up when you're sick—for example, when you have the flu or an infection. You'll need to take special care of yourself during these times. The guide that begins on page 33 can help you do this.

Signs of High Blood Glucose

Some common signs of high blood glucose are having a dry mouth, being thirsty, and urinating

often. Other signs include feeling tired, having blurred vision, and losing weight without trying. If your glucose is very high, you may have stomach pain, feel sick to your stomach, or even throw up. This is an emergency and you need to go to the hospital right away.



Frequent urination can be a sign of high blood sugar.

If you have any signs that your blood glucose is high, check your blood. In your logbook or on your record sheet, write down your glucose reading and the time you did the test. If your glucose is high, think about what could have caused it to go up. If you think you know of something, write this down beside your glucose reading.

Preventing High Blood Glucose

Keep a balance

Try to stay with your food and activity plan as much as you can. Drink water.
Take your diabetes medicine about the same time each day.
Work with your health care team to set goals for weight, blood glucose level, and activity.



Balance is the key to taking charge of your diabetes.

Test your blood glucose

Keep track of your blood glucose and go over your records often. You'll learn how certain foods or activities affect your glucose.

Show your records to your health care team. Ask how you can change your food, activity, and medicine to avoid or treat high blood glucose. Ask when you should call for help.

Taking Care of Yourself When You're Sick



You'll need to take special care of yourself when you're sick. The tips that follow can help you do this.

Keep Taking Medicine

Be sure to keep taking your diabetes pills or insulin. Don't stop taking them even if you can't eat. Your health care provider may even advise you to take more insulin during sickness.

Keep Eating

Try to eat the same amount of fruits and breads as usual. If you can, eat your regular diet. If you're having trouble doing this, use **carbohydrate choices or servings**: eat enough soft foods or drink enough liquids to take the place of the fruits and breads you usually eat.


What to Eat or Drink When You're Sick

Foods that have 15 grams carbohydrate or one carbohydrate serving

Food Item	Amount		
Fruit juice	¹ / ₂ cup		
Fruit-flavored drink	¹ / ₂ cup		
Soda pop (regular, not diet)	¹ / ₂ cup		
*JeII-O® (regular, not sugar-free)	¹ / ₂ cup		
*Popsicle® (regular, not sugar-free)	¹ / ₂ twin		
Sherbet	¹ / ₂ cup		
Saltine crackers	6 squares		
Bread	1 slice		
Milk	1 cup		
Soup	1 cup		
ce cream (regular)	¹ / ₂ cup		
Apple sauce	¹ / ₂ cup		
Pudding (regular)	¹ / ₄ cup		
Macaroni, noodles, rice,	1/3 cup (cooked)		
Potatoes, beans, cereal Suse of trade names is for identification only and does not imply endorsement by the U.S.	¹ / ₂ cup (cooked)		

Drink Liquids

Drink extra liquids. Try to drink at least ½ cup (4 ounces) to ¾ cup (6 ounces) every half-hour to hour, even if you have to do this in small sips. These liquids should not have calories. Water, diet soda pop, or tea without sugar are good choices.



Be sure to drink extra fluids when you're sick.

Check for Changes

- Weigh yourself every day. Losing weight without trying is a sign of high blood glucose.
- Check your temperature every morning and evening. A fever may be a sign of infection.
- Every 4 to 6 hours, check how you're breathing and decide how alert you feel. Having trouble breathing, feeling more sleepy than usual, or not thinking clearly can be danger signs.


Keep Records

Use the "Records for Sick Days," starting on page 79. Ask a family member or friend to help if you need it.

Call for Help

Ask your health care provider when you should call. During your sick times, you may need to call every day for advice.

You should call your health care provider or go to an emergency room if any of the following happens:




You may need to call your doctor daily when you're sick.

- You feel too sick to eat normally and for more than 6 hours can't keep food or liquids down.
- You have severe diarrhea (loose bowel movement).
- You lose 5 pounds or more without trying to.
- Your temperature is over 101°F.
- Your blood glucose level is lower than 60 mg/dL or stays over 300 mg/dL.
- You're having trouble breathing.
- You feel sleepy or can't think clearly.

Managing Your Diabetes at Work, School, and During Travel

Staying in charge of your diabetes no matter what your day holds—work, school, travel, or special events—takes planning ahead. Many days will go smoothly, but some days will hold surprises, such as extra activity or delays that throw your schedule off.

Plan ahead for these times by always keeping a treatment for low blood glucose with you (see page 26 for some choices). If you have any signs that your



Stay in charge of your diabetes—no matter what your day holds—by planning ahead.

glucose may be low (see page 25), go ahead and treat it right away.

Stay as close to your eating, activity, and medicine schedule as you can. Keep track of your blood glucose so you can pick up changes early. Always wear or carry identification that says you have diabetes.

Talk with your health care team about your planned schedule and activities. Ask for help in planning ahead for work, school, travel, and special events. When you read the rest of this section, you may think of more questions to ask.

At Work and School

Talk with your health care team about the type of activity you do at work or at school. From time to time, you and your health care team may need to make changes in your activity, medicine, or eating.

Many people take supplies for checking their glucose to school or work so they can check if at regular break times. Some people choose to show their fellow




Talk with your health care provider about balancing your daily activities and your glucose levels.

workers, their teachers, or their classmates how to help if they should ever have a problem. They teach them how to tell when their glucose is low and how to treat it (see pages 25–28). Some people like to have written steps on file at their place of work or with their teacher.

During Travel

When you plan a trip, think about your day-to-day schedule and try to stay as close to it as you can. For example, if you



Trips can hold surprises. Plan ahead for delays and changes.

usually check your blood glucose at noon and then eat lunch, plan to do this on your trip, as well. Trips can hold surprises—in delays and changes. Even the types of food and supplies you can buy on your trip may not be the same as those you get at home.

Before you travel, work with your health care provider to plan your timing for medicine, food, and activity. Talk about what to do if you find changes in your glucose readings.

Plan ahead for trips:

- Keep snacks with you that could be used to prevent—or treat—low blood glucose.
- Carry extra food and drink supplies with you, such as cracker packs and small cans of juices or bottled water.
- Carry blood glucose testing supplies with you.
- Take along all the diabetes medicine you'll need. Keep medicines in the original pharmacy container with the printed label that clearly identifies the medicine.

When you travel, be sure to

- Test your blood glucose often and keep track of it.
- Wear identification that says you have diabetes.
- Let others know how they can help you.
- Check new airline travel tips by contacting the Federal Aviation Administration (FAA) at http://www.faa.gov or 800-322-7873.

If you're traveling in a different time zone, you may need to change your timing of food, medicine, and activity. Ask your health care provider to help you with this. Talk about the food and drink choices that would be healthy for you. If you'll be in another country, ask your doctor to write a letter explaining that you have diabetes. It's also a good idea to get your doctor to write a prescription for you to get insulin or supplies if needed.



No matter where you travel, you can take charge of your diabetes.

4 Feelings About Having Diabetes

Living with diabetes isn't easy. It's normal to feel troubled about it. Tell your health care team how you feel. Point out any problems you have with your diabetes care plan. Your diabetes educator or other health care provider may be able to help you think of ways to deal with these problems.




Ask your family to help you manage your diabetes.

Talk about the stresses you feel at home, school, and work. How do you cope with these pressures? If your feelings are getting in the way of taking care of yourself, you need to ask for help.

Support Groups

It helps to talk with other people who have problems like your own. You may want to think about joining a diabetes **support group**. In support groups, people who have just found out they have diabetes can learn from people who have lived with it for a long time. People can talk about and share how they deal with their diabetes. They can also talk about how they take care of their health, how they prepare food, and how they get physical activity. Family members who do not have diabetes may want to join a support group, too. Ask your health care team about support groups for people with diabetes and their families and friends.



It can help to talk with other people who have problems like your own.

If there is not a support group in your area, you may want to call a diabetes organization (see the list on pages 127–129) about starting a group.

Counseling


One-on-one and family counseling sessions may also help. Be sure to see a counselor who knows about diabetes and its care. Ask your health care provider to help you find a counselor.

5 Eye Problems

Diabetic eye disease (also called diabetic **retinopathy**) is a serious problem that can lead to loss of sight. There's a lot you can do to take charge and prevent such problems. Research shows that keeping your **blood glucose** level closer to normal can prevent or delay the onset of diabetic eye disease. Keeping your blood pressure under control is also important. Finding and treating eye problems early can help save sight.

Signs of Diabetic Eye Disease

Since diabetic eye disease may be developing even when your sight is good, regular **dilated** eye exams are important for finding problems early. Some people may notice signs of vision changes. If you're having trouble reading, if



If you're having trouble with your vision, talk with your health care team or eye doctor.

your vision is blurred, or if you're seeing rings around lights, dark spots, or flashing lights, you may have eye problems. Be sure to tell your health care team or eye doctor about any eye problems you may have.

Protecting Your Sight

Keep Your Blood Glucose Under Control

High blood glucose can damage your eyes as time goes by. Work with your health care team to keep your blood glucose levels in the target range.

Keep Your Blood Pressure Under Control


High blood pressure can damage your eyes. Have your health care provider check your **blood pressure** at least 4 times a year. If your blood pressure is higher than 130/80, ask your health care provider how to keep your blood pressure at a healthy level. You may need medicine to keep your blood pressure at a healthy level.



You may want to check your blood pressure at home.

Get Regular Eye Exams

Even if you're seeing fine, you need regular, complete dilated eye exams to protect your sight. Ask your health care provider to help you find an eye doctor who cares for people with diabetes. Before



Get a complete eye exam each year.

the exam, a doctor or nurse will put drops in your eyes to dilate the pupils.


You should have your eyes **dilated** and examined at least once a year. Keep track of these exams by using the record sheets starting on page 101. Even if you've lost your sight from diabetic eye disease, you still need to have regular eye care. If you haven't already had a complete eye exam, you should have one now if any of these conditions apply to you:

- You've had **type 1 diabetes** for 5 or more years.
- You have **type 2 diabetes**.
- You're going through puberty and you have diabetes.
- You're pregnant and you have diabetes.
- You're planning to become pregnant and you have diabetes.

If you can't afford an eye exam, ask about a payment plan or a free exam. If you're 65 or older, Medicare may pay for diabetic eye exams (but not glasses). Ask your eye doctor to accept the Medicare fee as full payment.

Discuss Your Physical Activity Plan

If you have diabetic eye disease, talk with your health care provider about the kind of physical activity that is best for you.



Think of a way to remind yourself to get an eye exam each year.

Treating Diabetic Eye Disease

Treating eye problems early can help save sight. **Laser surgery** may help people who have advanced diabetic eye disease. An operation called a **vitrectomy** may help those who have lost their sight from bleeding in the back of the eye.


If your sight is poor, an eye doctor who is an expert in low vision may be able to give you glasses or other devices that can help you use your limited vision more fully. You may want to ask your health care provider about support groups and job training for people with poor vision.

6 Kidney Problems

Diabetes can cause **diabetic kidney disease** (also called diabetic **nephropathy**), which can lead to kidney failure. There's a lot you can do to take charge and prevent kidney problems. A recent study shows that controlling your **blood glucose** can prevent or delay the onset of kidney disease. Keeping your blood pressure under control is also important.

The **kidneys** keep the right amount of water in the body and help filter out harmful wastes. These wastes, called **urea**, then pass from the body in the urine. Diabetes can cause kidney disease by damaging the parts of the kidneys that filter out wastes. When the kidneys fail, a person has to have his or her blood filtered through a machine (a treatment called **dialysis**) several times a week or has to get a kidney transplant.

Take care of your kidneys by keeping your blood glucose and blood pressure in balance.



Testing Your Kidneys

Your health care provider can learn how well your kidneys are working by testing for **microalbumin** (a protein) in the urine. Microalbumin in the urine is an early sign of diabetic kidney disease. You

should have your urine checked for microalbumin every year.

Your health care provider can also do a yearly blood test to measure your kidney function. If the tests show microalbumin in the urine or if your kidney function isn't normal, you'll need to be checked more often.



Work with your health care provider to prevent or treat kidney problems.

Starting on page 101, write down the dates and the results of these tests. Ask your health care provider to explain what the results mean.

Protecting Your Kidneys


Keep Your Blood Glucose Under Control

High blood glucose can damage your kidneys as time goes by. Work with your health care team to keep your glucose levels as close to normal as you can.

Keep Your Blood Pressure In Balance

High blood pressure (or hypertension) can

damage your kidneys. You may want to check your blood pressure at home to be sure it stays lower than 130/80. Have your health care provider check your blood pressure at least 4 times a year. Your doctor may have you take a blood pressure pill, called an ACE inhibitor, to help protect your kidneys.



Controlling your blood glucose levels and your blood pressure may help to prevent or delay kidney failure.

To Lower Your Blood Pressure:

- 1. Practice these steps:
 - Maintain a healthy weight.
 - Be acive every day.
 - Eat fewer foods high in salt and sodium.
 - Eat more fruits and vegetables, whole grain breads and cereals, and lowfat dairy products.
- 2. Take your medicine the way your doctor tells you.
- 3. Have your blood pressure checked often.

Call your health care provider right away if you have any of these signs of kidney infections:

- Back pain.
- Chills.
- Fever.
- **Ketones** in the urine.



Tell your health care provider if you have any signs of kidney or bladder infection.

Your health care provider will test your urine. If you have a bladder or kidney infection, you'll be given medicine to stop the infection. After you take all the medicine, have your urine checked again to be sure the infection is gone.

Know the Effects of Some Medicines and X-Ray Dyes

If you have kidney disease, ask your health care provider about the possible effects that some medicines and X-ray dyes can have on your kidneys.

7 Heart and Blood Vessel Problems

Heart and blood vessel problems are the main causes of sickness and death among people with diabetes. These problems can lead to **high blood pressure**, **heart attacks**, and **strokes**. Heart and blood vessel problems can also cause poor circulation (blood flow) in the legs and feet.

You're more likely to have heart and blood vessel problems if you smoke cigarettes, have high blood pressure, or have too much **cholesterol** or other fats in your blood. Talk with your health care team about what you can do to lower your risk for heart and blood vessel problems. Ask about taking a daily aspirin to help prevent heart and blood vessel problems.

You can do a lot to keep your heart and blood vessels healthy.

Signs of Heart and Blood Vessel Problems

If you feel dizzy, have sudden loss of sight, slur your speech, or feel numb or weak in one arm or leg, you may be having serious heart and blood vessel problems. Your blood may not be getting to your brain as well as it should.


Danger signs of circulation problems to the heart include chest pain or pressure, shortness of breath, swollen ankles, or irregular heartbeats. If you have any of these signs, go to an emergency room or call your health care provider right away.

Signs of circulation problems to your legs are pain or cramping in your buttocks, thighs, or calves during physical activity. Even if this pain goes away with rest, report it to your health care provider.

Preventing and Controlling Heart and Blood Vessel Problems

Eat Right and Get Physical Activity

Choose a healthy diet, low in salt. Work with a dietitian to plan healthy meals. If you're overweight, talk about how to safely lose weight. Ask about a physical activity or exercise program for you. See pages 14–18 to read more about healthy choices for food and physical activity.



If you're overweight, talk with your dietitian about how to safely lose weight.

Don't Use Tobacco

Smoking cigarettes causes hundreds of thousands of deaths each year. When you have diabetes and also use tobacco, the risk of heart and blood vessel problems is even greater. One of the best choices you

can make for your health is to never start smoking—or if you smoke, to quit.

At least once a year, your health care provider will ask you about tobacco use. If you smoke, talk to your provider about ways to help you stop.



Not smoking is the healthiest choice you'll make for your heart.


Check Your Blood Pressure

Get your **blood pressure** checked at each visit. Record these numbers on the record sheets starting on page 91. If your blood pressure is higher than 130/80, ask what steps to take to reach your goal.

If your blood pressure is still high after 3 months, you may need medicine to help control it. Many medicines are available to treat high blood pressure. If you have side effects from the medicine, ask your health care provider to change it. Talk to your health care team about whether you need medicine to take charge of your blood pressure.

Check Your Cholesterol

Get your cholesterol checked once a year. Record the results on page 101. Your total cholesterol should be lower than 200 mg/dL (milligrams per deciliter). Ask your health care team to explain what your **HDL** and **LDL** levels are.



Choose heart-healthy foods for your meal plan.

If your cholesterol is higher than 200 mg/dL on two or more checks, you can do several things to lower it. You can work with your health care team to improve your **blood glucose** control, you can lose weight (if you're overweight), and you can cut down on foods that are high in fat and cholesterol. Ask your health care team about foods that are low in fats. Also ask about a physical activity program.

Ask your health care provider what steps to take to reach your LDL cholesterol goal. You may need a medicine to help control it. Ask if you need aspirin to prevent heart attack or stroke.

Ask If You Need an Electrocardiogram (EKG)

If you're having heart and blood circulation problems, an **EKG** may help you and your health care provider know if you need to change your treatment.

8 Nerve Damage

Diabetic nerve damage (also called diabetic **neuropathy**) is a problem for many people with diabetes. Over time, **high blood glucose** levels damage the delicate coating of nerves. This damage can cause many problems, such as pain in your feet. There's a lot you can do to take charge and prevent nerve damage. A recent study shows that controlling your **blood glucose** can help prevent or delay these problems. Controlling your blood glucose may also help reduce the pain from some types of nerve damage.

Some Signs of Diabetic Nerve Damage

Some signs of diabetic nerve damage are pain, burning, tingling, or loss of feeling in the feet and hands. It can cause you to sweat abnormally, make it hard for you to tell when your blood glucose is low, and make you feel light-headed when you stand up.

Nerve damage can lead to other problems. Some people develop problems swallowing and keeping



Having trouble telling your glucose is low may be a sign of nerve damage.

food down. Nerve damage can also cause bowel problems, make it hard to urinate, cause dribbling



Tell your health care provider if you have trouble with sexual function.

with urination, and lead to bladder and kidney infections. Many people with nerve damage have trouble having sex. For example, men can have trouble keeping their penis erect, a problem called **impotence** (erectile dysfunction). If you have any of these problems, tell your health care provider. There are ways to help

in many cases.

Protecting Your Nerves from Damage

Keep Your Blood Glucose in Control


High blood glucose can damage your nerves as time goes by. Work with your health care team to keep your glucose levels as close to normal as you can.

Have a Physical Activity Plan

Physical activity or exercise may help keep some nerves healthy, such as those in your feet. Ask your health care team about an activity that is healthy for you.

Get Tested for Nerve Damage

Nerve damage can happen slowly. You may not even be aware you're losing feeling in your feet. Ask your health care provider to check your feet at each visit. At least once a year, your provider should test how well you can sense



At least once a year, your health care provider should do a complete check of your feet and nerves.

temperature, pinprick, vibration, and position in your feet. If you have signs of nerve damage, your provider may want to do more tests. Testing can help your provider know what is wrong and how to treat it. Keep track of your foot exams on the record sheets starting on page 101.

For more information on foot care, call the National Diabetes Information Clearinghouse at 1-800-860-8747.

Check Your Feet for Changes

If you've lost feeling in your feet, you'll need to take special care of them. Check your feet each day. Wear shoes that fit well. You'll read more about foot care in the next chapter.

9 Foot Problems

Nerve damage, circulation problems, and infections can cause serious foot problems for people with diabetes. There's a lot you can do to prevent problems with your feet. Controlling your **blood glucose** and not smoking or using tobacco can help protect your feet. You can



Take extra care of your feet to prevent injuries.

also take some simple safeguards each day to care for and protect your feet. Over half of diabetesrelated amputations can be prevented with regular exams and patient education.

It's helpful to understand why foot problems happen. Nerve damage can cause you to lose feeling in your feet. Sometimes nerve damage can deform or misshape your feet, causing pressure points that can turn into blisters, sores, or **ulcers**. Poor circulation can make these injuries slow to heal.

Signs of Foot Problems

Your feet may tingle, burn, or hurt. You may not be able to feel touch, heat, or cold very well. The shape of your feet can change over time. There may even be changes in the color and temperature of your feet. Some people lose hair on their toes, feet, and lower legs. The skin on your feet may be dry and cracked. Toenails may turn thick and yellow. Fungus

infections can grow between your toes. Blisters, sores, ulcers, infected **corns**, and ingrown toenails need to be seen by your health care provider or foot doctor (podiatrist) right away.

Protecting Your Feet

Get Your Health Care Provider to Check Your Feet at Least 4 Times a Year

Ask your health care provider to look at your feet at least 4 times a year. As a reminder, take off your shoes and socks when you're in the exam room. Have your sense of feeling and your pulses checked at least once a year. If you have nerve damage, deformed or misshaped feet, or a circulation problem, your feet need special care. Ask your health care provider to show you how to care for your feet. Also ask if special shoes would help you.



Ask your health care provider to check your feet at least 4 times a year.

Check Your Feet Each Day

You may have serious foot problems yet feel no pain. Look at your feet every day to see if you have scratches, cracks, cuts, or blisters. Always check between your toes and on the bottoms of your feet. If you can't bend over to see the bottoms of your feet, use a mirror that won't break. If you can't see well, ask a family member or friend to help you. Call your health care provider at once if you have a sore on your foot. Sores can get worse quickly.

Wash Your Feet Daily

Wash your feet every day. Dry them with care, especially between the toes. Don't soak your feet—it can dry out your skin, and dry skin can lead to infections. Rub lotion or cream on the tops and bottoms of your feet—but not between your toes. Moisture between the toes will let germs grow that could cause an infection. Ask your health care provider for the name of a good lotion or cream.



Be sure to dry between your toes.

Trim Your Toenails Carefully

Trim your toenails after you've washed and dried your feet—the nails will be softer and safer to cut. Trim the nails to follow the natural curve of your toes. Don't cut into the corners. Use an emery board to smooth the edges.

If you can't see well, or if your nails are thick or yellowed, get them trimmed by a foot doctor or another health care provider. Ask your health care provider for the name of a foot doctor. If you see redness around the nails, see your health care provider at once.

Treat Corns and Calluses Gently

Don't cut **corns** and **calluses**. Ask your health care provider how to gently use a **pumice stone** to rub them. Don't use razor blades, corn plasters, or liquid corn or callus removers—they can damage your skin.

Protect Your Feet from Heat and Cold

Hot water or hot surfaces are a danger to your feet. Before bathing, test the water with a bath thermometer (90° to 95°F is safe) or with your elbow. Wear shoes and socks when you walk on hot surfaces, such as beaches or the pavement around swimming pools. In summer, be sure to use sunscreen on the tops of your feet.



Wear shoes to protect your feet from hot surfaces.

You also need to protect your feet from the cold. In winter, wear socks and footwear such as fleecelined boots to protect your feet. If your feet are cold at night, wear socks. Don't use hot water bottles, heating pads, or electric blankets—they can burn your feet. Don't use strong antiseptic solutions or adhesive tape on your feet.

Always Wear Shoes and Socks

Wear shoes and socks at all times. Don't walk barefoot—not even indoors.



Wear shoes that fit well and protect your feet.

Wear shoes that fit well and protect your feet. Don't

wear shoes that have plastic uppers, and don't wear sandals with thongs between the toes. Ask your health care provider what types of shoes are good choices for you.

New shoes should be comfortable at the time you buy them—don't expect them to stretch out. Slowly break in new shoes by wearing them only 1 or 2 hours a day.

Always wear socks or stockings with your shoes. Choose socks made of cotton or wool—they help keep your feet dry.

Before you put on your shoes each time, look and feel inside them. Check for any loose objects, nail points, torn linings, and rough areas—these can cause injuries. If your shoes aren't smooth inside, wear other shoes.

Be Physically Active

Physical activity can help increase the circulation in your feet. There are many ways you can exercise your feet, even during times you're not able to walk. Ask your health care team about things you can do to exercise your feet and legs.

For more information on foot care, call the National Diabetes Information Clearinghouse at 1-800-860-8747.



Being active is a healthy way to live.

10 Dental Disease



Healthy teeth and gums depend on regular care and controlling your blood glucose levels.

Because of high blood glucose, people with diabetes are more likely to have problems with their teeth and gums. There's a lot you can do to take charge and prevent these problems. Caring for your teeth and gums every day can help keep them healthy. Keeping your blood glucose under control is also important. Regular, complete dental care helps prevent dental disease.

Signs of Dental Disease

Sore, swollen, and red gums that bleed when you brush your teeth are a sign of a dental problem called **gingivitis**. Another problem, called **periodontitis**, happens when your gums shrink or pull away from your teeth. Like all infections, dental infections can make your blood glucose go up.

Preventing Dental Problems

Keep Your Blood Glucose Under Control

High blood glucose can cause problems with your teeth and gums. Work with your health care team to keep your glucose levels as close to normal as you can.

Brush Your Teeth Often

Brush your teeth at least twice a day to prevent gum disease and tooth loss. Be sure to brush before you go to sleep. Use a soft toothbrush and toothpaste with fluoride. To help keep bacteria from growing on your toothbrush, rinse it after each brushing and store it upright with the bristles at the top. Get a new toothbrush at least every 3 months.



Protect your teeth by brushing twice or more a day and flossing each day.

Floss Your Teeth Daily

Besides brushing, you need to floss between your teeth each day to help remove **plaque**, a film that forms on teeth and can cause tooth problems. Flossing also helps keep your gums healthy. Your dentist or dental hygienist will help you choose a good method to remove plaque, such as dental floss, bridge cleaners, or water spray. If you're not sure of the right way to brush or floss, ask your dentist or dental hygienist for help.

Get Regular Dental Care

Get your teeth cleaned and checked at your dentist's office at least once every 6 months. If you don't have a dentist, find one or ask your health care provider for the name of a dentist in your community.



See your dentist at least once every 6 months.

See your dentist right away if you have trouble

chewing or any signs of dental disease, including bad breath, a bad taste in your mouth, bleeding or sore gums, red or swollen gums, or sore or loose teeth.

Give your dentist the name and telephone number of your diabetes health care provider. Each time you visit, remind your dentist that you have diabetes.

Plan dental visits so they don't change the times you take your **insulin** and meals. Don't skip a meal or diabetes medicine before your visit. Right after breakfast may be a good time for your visit.

11 Vaccinations

If you have diabetes, take extra care to keep up-to-date on your **vaccinations** (also called **immunizations**). Vaccines can prevent illnesses that can be very serious for people with diabetes. This section talks about some vaccines you need to know about.

Influenza Vaccine

Influenza (often called the **flu**) is not just a bad cold. It's a serious illness that can lead to pneumonia and even death. The flu spreads when influenza viruses pass from one person to the nose or throat of others. Signs of the flu may include sudden high fever, chills, body aches, sore throat, runny nose, dry cough, and headache.



The flu is a serious illness that can put you in the hospital. A yearly flu shot can help prevent this.

People with diabetes who come down with the flu may become very sick and may even have to go to a hospital. If you get the flu, you'll need to take special care of yourself (see pages 33–36).

You can help keep yourself from getting the flu by getting a flu shot every year. Everyone with diabetes—even pregnant women—should get a yearly flu shot. The best time to get one is between October and mid-November, before the flu season begins. This vaccine is fully covered under Medicare Part B.

Pneumococcal Vaccine

Pneumococcal disease is a major source of illness and death. It can cause serious infections of the lungs (pneumonia), the blood (bacteremia), and the covering of the brain (meningitis). Pneumococcal polysaccharide vaccine (often called PPV) can help prevent this disease.

PPV can be given at the same time as the flu vaccine—or at any time of the year. Most people only have to take PPV once in their life. Ask your health care provider whether you might need a second vaccination. This vaccine is fully covered under Medicare Part B.

Tetanus/Diphtheria (Td) Toxoid

Tetanus (or lockjaw) and **diphtheria** are serious diseases. Tetanus is caused by a germ that enters the body through a cut or wound. Diphtheria spreads when germs pass from one person to the nose or throat of others.

You can help prevent tetanus and diphtheria with a combined shot called Td toxoid. Most people get Td toxoid as part of their routine childhood vaccinations, but all adults need a Td booster shot every 10 years. Other vaccines may be given at the same time as Td toxoid.

Other Vaccines


You may need vaccines to protect you against other illnesses. Ask your health care provider if you need any of these:

- Measles/Mumps/Rubella vaccine.
- Hepatitis A and B vaccines.
- Varicella (chicken pox) vaccine.
- Polio vaccine.
- Vaccines for travel to other countries.

How to Get More Information

Call the immunization program in your state health department to find out where you can get vaccinations in your area. Keep your vaccination records up-to-date so you and your health care provider will know what vaccines you may need. You can record this information on the record sheets starting on page 101 of this book.

For more information on vaccination, call the CDC National Immunization Hotline at 1-800-CDC-INFO (English and Spanish). This is a toll-free call.



12 Pregnancy and Women's Health

Becoming Pregnant When You Have Diabetes

Women with diabetes can have healthy babies, but it takes planning ahead and effort. Pregnancy can make both high and low blood glucose levels happen more often. It can make diabetic eye disease and diabetic kidney disease worse. High glucose levels during pregnancy are dangerous for the baby, too.

If you don't want to become pregnant, talk with your health care provider about birth control.



You can protect your baby and yourself by controlling your blood glucose before and during pregnancy.

Protecting Your Baby and Yourself

Keeping your glucose levels near normal before and during pregnancy can help protect you and your baby. That's why it's so important to plan your pregnancies ahead of time.



Your blood glucose and A1C records will help you and your health care team know when your glucose range is safe for pregnancy.

If you want to have a baby, discuss it with your health care provider. Work with your diabetes care team to get and keep your blood glucose in the normal or near-normal range before you become pregnant. Your glucose records and your **A1C** test results will show when you have maintained a safe range for a period of time.

You may need to change your **meal plan** and your usual physical activity, and you may need to take more frequent **insulin** shots. Testing your glucose several times a day will help you see how well you're balancing things. Record the test results in your logbook or on a log sheet (see sample pages on 107–108).

Get a complete check of your eyes and **kidneys** before you try to become pregnant. Don't smoke, drink alcohol, or use drugs—doing these things can harm you and your baby.

All women who could become pregnant should take folic acid (400 micrograms) every day. An easy way to be sure you're getting enough folic acid is to take a vitamin with folic acid in it.

Think about breast feeding your baby. Breast feeding has many benefits for you and your baby.

Having Diabetes During Pregnancy

Some women have diabetes only when they're pregnant. This condition, which is called **gestational diabetes**, can be controlled just like other kinds of diabetes. Glucose control is the key. Your health care team can help you take charge of gestational diabetes. You are more likely to develop type 2 diabetes. Check again for diabetes at least 6 weeks after your baby is born and at regular times for the rest of your life.



If you learn you have diabetes when you're pregnant, work closely with your health care team to stay healthy.

Controlling Diabetes for Women's Health

Some women with diabetes may have special problems, such as **bladder** infections. See pages 49–50 to find out about the signs of bladder and kidney infections. If you have an infection, it needs to be treated right away. Call your doctor.

Some women get **yeast infections** in their **vagina**, especially when their blood glucose is high. A sign of a yeast infection may be itching in the vagina. If you notice vaginal itching, tell your health care provider, who can tell you about medicines you can buy at the drugstore and about how to prevent yeast infections.

Some women with diabetes may have trouble with sexual function. Discomfort caused by vaginal itching or dryness can be treated.

Ask your doctor how often you should get a Pap smear and a mammogram (breast X-ray). Regular Pap smears and mammograms help detect cervical and breast cancer early. All women—whether or not they have diabetes—need to have these tests regularly.



Getting Pap smears and mammograms is important to every woman's health.

RECORDS

Records for Sick Days
Checks and Goals for Each Visit 89
Checks and Goals for Each Year 99
Glucose Record Sheets
Your Health Care Team 109

Records for Sick Days

How often	Question	Answ	er
Every day	How much do you weigh today?		pounds
Every evening	How much liquid did you drink today?		glasses
Every morning and every evening	What is your temperature?		
Every 4 hours or before every meal	How much insulin did you take?	Time	Dose
Every 4 hours	What is your blood glucose level?	Time	Blood glucose
Every 4 hours or each time you pass urine	What are your urine ketones?	Time	Ketones

Every 4 to 6 hours	How are you breathing?	Time Condition
o riours	broatimig.	

Reminders for Sick Days

Call your health care provider if any of these happen to you:

- You feel too sick to eat normally and are unable to keep down food for more than 6 hours.
- You're having severe diarrhea.
- You lose 5 pounds or more.
- Your temperature is over 101°F.
- Your blood glucose is lower than 60 mg/dL or remains over 300 mg/dL.
- You have moderate or large amounts of ketones in your urine.
- You're having trouble breathing.
- You feel sleepy or can't think clearly.

If you feel sleepy or can't think clearly, have someone call your health care provider or take you to an emergency room.

Records for Sick Days

How often	Question	Answer
Every day	How much do you weigh today?	pounds
Every evening	How much liquid did you drink today?	glasses
Every morning and every evening	What is your temperature?	a.m. p.m.
Every 4 hours or before every meal	How much insulin did you take?	Time Dose
Every 4 hours	What is your blood glucose level?	Blood Time glucose
Every 4 hours or each time you pass urine	What are your urine ketones?	Time Ketones

81

Every 4 to 6 hours	How are you	Time Condition
0 110ul 3	breathing?	

Reminders for Sick Days

Call your health care provider if any of these happen to you:

- You feel too sick to eat normally and are unable to keep down food for more than 6 hours.
- You're having severe diarrhea.
- You lose 5 pounds or more.
- Your temperature is over 101°F.
- Your blood glucose is lower than 60 mg/dL or remains over 300 mg/dL.
- You have moderate or large amounts of ketones in your urine.
- You're having trouble breathing.
- You feel sleepy or can't think clearly.

If you feel sleepy or can't think clearly, have someone call your health care provider or take you to an emergency room.

Records for Sick Days

How often	Question	Answe	er
Every day	How much do you weigh today?		pounds
Every evening	How much liquid did you drink today?		glasses
Every morning and every evening	What is your temperature?		
Every 4 hours or before every meal	How much insulin did you take?	Time	Dose
Every 4 hours	What is your blood glucose level?	Time	Blood glucose
Every 4 hours or each time you pass urine	What are your urine ketones?	Time	Ketones



Every 4 to 6 hours	How are you	Time Condition
0 Hours	breathing?	

Reminders for Sick Days

Call your health care provider if any of these happen to you:

- You feel too sick to eat normally and are unable to keep down food for more than 6 hours.
- You're having severe diarrhea.
- You lose 5 pounds or more.
- Your temperature is over 101°F.
- Your blood glucose is lower than 60 mg/dL or remains over 300 mg/dL.
- You have moderate or large amounts of ketones in your urine.
- You're having trouble breathing.
- You feel sleepy or can't think clearly.

If you feel sleepy or can't think clearly, have someone call your health care provider or take you to an emergency room.

Records for Sick Days

How often	Question	Answe	er
Every day	How much do you weigh today?		pounds
Every evening	How much liquid did you drink today?		glasses
Every morning and every evening	What is your temperature?		
Every 4 hours or before every meal	How much insulin did you take?	Time	Dose
Every 4 hours	What is your blood glucose level?	Time	Blood glucose
Every 4 hours or each time you pass urine	What are your urine ketones?	Time	Ketones



Every 4 to 6 hours	How are you breathing?	Time Condition
OTIONIS	breatining.	

Reminders for Sick Days

Call your health care provider if any of these happen to you:

- You feel too sick to eat normally and are unable to keep down food for more than 6 hours.
- You're having severe diarrhea.
- You lose 5 pounds or more.
- Your temperature is over 101°F.
- Your blood glucose is lower than 60 mg/dL or remains over 300 mg/dL.
- You have moderate or large amounts of ketones in your urine.
- You're having trouble breathing.
- You feel sleepy or can't think clearly.

If you feel sleepy or can't think clearly, have someone call your health care provider or take you to an emergency room.

Records for Sick Days

How often	Question	Answer
Every day	How much do you weigh today?	pounds
Every evening	How much liquid did you drink today?	glasses
Every morning and every evening	What is your temperature?	a.m. p.m.
Every 4 hours or before every meal	How much insulin did you take?	Time Dose
Every 4 hours	What is your blood glucose level?	Blood Time glucose
Every 4 hours or each time you pass urine	What are your urine ketones?	Time Ketones



Every 4 to 6 hours	How are you breathing?	Time (Condition
o riours	broatimig.		

Reminders for Sick Days

Call your health care provider if any of these happen to you:

- You feel too sick to eat normally and are unable to keep down food for more than 6 hours.
- You're having severe diarrhea.
- You lose 5 pounds or more.
- Your temperature is over 101°F.
- Your blood glucose is lower than 60 mg/dL or remains over 300 mg/dL.
- You have moderate or large amounts of ketones in your urine.
- You're having trouble breathing.
- You feel sleepy or can't think clearly.

If you feel sleepy or can't think clearly, have someone call your health care provider or take you to an emergency room.

Things to Do at Each Visit with Your Health Care Provider

- Bring your blood glucose record book and go over the readings with your provider.
- Get an A1C test every 3 months. Write down the result and set a target goal for your next test. (See the sample form on page 90.)
- Get your weight checked and write it down. You may want to set a goal for your next visit.
- Get your blood pressure checked and write it down. You may want to set a goal for your next visit.
- Get your feet checked at every visit as needed.
- Bring a list of questions or other things you want to talk about.
- Bring your reminder sheet about "Things to Do at Least Once a Year" (see page 99) to help keep track of these.

Each Visit

Have your health care provider do these tests and set goals with you. (Record dates and results in the boxes below.)

Tests and Goals			Dates an	Dates and Results		
Blood Glucose (mg/dL)	2/1/2000	0/11/5000	0007/87/6	1/2/5001	4/3/2001	
	145	118	180	105	110	
A1C	0.6	8.9	8.4	not done	8.2	
Test/Goal (%)	8,0	8.0	7.5		7.5	
Weight/Goal	180	9,11	7.11	021	991	
(spunod)	770	165	165	165	760	
Blood Pressure (goal: 120 180 mm Hg)	06/041	140 / 86	138 / 84	78/9£1	124 / 80	
Foot Check	>	,	`	'	/	

Cut here if you want to take only this page to visits with your health care provider.)

Each Visit

Have your health care provider do these tests and set goals with you. (Record dates and results in the boxes below.)

Tests and Goals		Dates an	Dates and Results	
Blood Glucose (mg/dL)				
A1C Test/Goal (%)				
Weight/Goal (pounds)				
Blood Pressure (goal:/mm Hg)				
Foot Check				

Each Visit

Have your health care provider do these tests and set goals with you. (Record dates and results in the boxes below.)

Tests and Goals		Dates an	Dates and Results	
Blood Glucose (mg/dL)				
A1C Test/Goal (%)				
Weight/Goal (pounds)				
Blood Pressure (goal:/_mm Hg)				
Foot Check				

Each Visit

Have your health care provider do these tests and set goals with you. (Record dates and results in the boxes below.)

Tests and Goals		Dates an	Dates and Results	
Blood Glucose (mg/dL)				
A1C Test/Goal (%)				
Weight/Goal (pounds)				
Blood Pressure (goal:/_mm Hg)				
Foot Check				

Each Visit

Have your health care provider do these tests and set goals with you. (Record dates and results in the boxes below.)

Tests and Goals		Dates an	Dates and Results	
Blood Glucose (mg/dL)				
A1C Test/Goal (%)				
Weight/Goal (pounds)				
Blood Pressure (goal:/_mm Hg)				
Foot Check				

Each Visit

Have your health care provider do these tests and set goals with you. (Record dates and results in the boxes below.)

Tests and Goals		Dates an	Dates and Results	
Blood Glucose (mg/dL)				
A1C Test/Goal (%)				
Weight/Goal (pounds)				
Blood Pressure (goal:/_mm Hg)				
Foot Check				

95

Each Visit

Have your health care provider do these tests and set goals with you. (Record dates and results in the boxes below.)

Tests and Goals		Dates an	Dates and Results	
Blood Glucose (mg/dL)				
A1C Test/Goal (%)				
Weight/Goal (pounds)				
Blood Pressure (goal:/_mm Hg)				
Foot Check				

Each Visit

Have your health care provider do these tests and set goals with you. (Record dates and results in the boxes below.)

Tests and Goals		Dates an	Dates and Results	
Blood Glucose (mg/dL)				
A1C Test/Goal (%)				
Weight/Goal (pounds)				
Blood Pressure (goal:/_mm Hg)				
Foot Check				

Cut here if you want to take only this page to visits with your health care provider.)

Each Visit

Have your health care provider do these tests and set goals with you. (Record dates and results in the boxes below.)

Tests and Goals		Dates an	Dates and Results	
Blood Glucose (mg/dL)				
A1C Test/Goal (%)				
Weight/Goal (pounds)				
Blood Pressure (goal:/mm Hg)				
Foot Check				

Things to Do At Least Once a Year

- Get a flu shot (October to mid-November).
- Get a pneumonia shot (if you've never had one) and every evvery 5 years.
- Get a dilated eye exam.
- Get a foot exam (including check of circulation and nerves).
- Get a kidney test.
 - Have your urine tested for microalbumin.
 - Have your blood tested for chemicals that measure your kidney function.
 - Get a 24-hour urine test (if your doctor advises).
- Get your blood fats checked for
 - Total cholesterol.
 - High-density lipoprotein (HDL).
 - Low-density lipoprotein (LDL).
 - Triglycerides.
- Get a dental exam (at least twice a year).
- Talk with your health care team about
 - How well you can tell when you have low blood glucose.
 - How you are treating high blood glucose.
 - Tobacco use (cigarettes, cigars, pipes, smokeless tobacco).
 - Your feelings about having diabetes.
 - Your plans for pregnancy (if a woman).
 - Other

Cut here if you want to take only this page to visits with your health care provider.)

At Least Once a Year

Have your health care provider do these tests and other services for you. (Record the dates and results in the boxes below.) You may want to set some goals for these.

Tests and Other Services			Õ	Dates and Results	Results		
Flu Shot	10/2/1999	10/2/1999 10/20/2000 11/1/2001	11/1/2001				
Urine Protein or Wicroalbumin (mg)	10/2/1999	10/2/1999 10/20/2000 11/1/2001 40 50 55	11/1/2001				
Blood Creatinine (mg/dL)	0.7	1.2	1.1				
Total Cholesterol (mg/dL)	061		175				
HDL Cholesterol (mg/dL)	30	35	40				
LDL Cholesterol (mg/dL)	051	140	135				
Triglycerides (mg/dL)	338	300	250	5			
Tobacco Use	5 cigars a day	2 cigares	0				
Eye Exam (dilated)	6661/11/8	8/11/1999 10/1/2000 10/20/2001	10/20/2001				
Foot Exam	10/2/1999	10/2/1999 10/20/2000 11/1/2001	11/1/2001				

(Cut here if you want to take only this page to visits with your health care provider.)

At Least Once a Year

Have your health care provider do these tests and other services for you. You may want to set some goals for these.

(Record the dates and results in the boxes below.)

Dates and Results Tests and Other Services Blood Creatinine (mg/dL) Total Cholesterol (mg/dL) HDL Cholesterol (mg/dL) LDL Cholesterol (mg/dL) Triglycerides (mg/dL) Microalbumin (mg) Eye Exam (dilated) Urine Protein or Tobacco Use Foot Exam Flu Shot

At Least Once a Year

Have your health care provider do these tests and other services for you. (Record the dates and results in the boxes below.) You may want to set some goals for these.

Tests and Other Services	Dates and Results
Flu Shot	
Urine Protein or Microalbumin (mg)	
Blood Creatinine (mg/dL)	
Total Cholesterol (mg/dL)	
HDL Cholesterol (mg/dL)	
LDL Cholesterol (mg/dL)	
Triglycerides (mg/dL)	
Tobacco Use	
Eye Exam (dilated)	
Foot Exam	

(Cut here if you want to take only this page to visits with your health care provider.)

(Cut here if you want to take only this page to visits with your health care provider.)

At Least Once a Year

Have your health care provider do these tests and other services for you. You may want to set some goals for these.

(Record the dates and results in the boxes below.)

Tests and Other Services	٥	Dates and Results	Results		
Flu Shot					
Urine Protein or Microalbumin (mg)					
Blood Creatinine (mg/dL)					
Total Cholesterol (mg/dL)					
HDL Cholesterol (mg/dL)					
LDL Cholesterol (mg/dL)					
Triglycerides (mg/dL)					
Tobacco Use					
Eye Exam (dilated)					
Foot Exam					

At Least Once a Year

Have your health care provider do these tests and other services for you. (Record the dates and results in the boxes below.) You may want to set some goals for these.

Tests and Other Services	Dates and Results
Flu Shot	
Urine Protein or Microalbumin (mg)	
Blood Creatinine (mg/dL)	
Total Cholesterol (mg/dL)	
HDL Cholesterol (mg/dL)	
LDL Cholesterol (mg/dL)	
Triglycerides (mg/dL)	
Tobacco Use	
Eye Exam (dilated)	
Foot Exam	

Cut here if you want to take only this page to visits with your health care provider.)

Glucose Log Sheet for People Who Do Not Use Insulin Personal target Goal: Fasting 90-130/After meals less then 180

Medicine: Glyburide 10 mg twice a day. glucophage 1000 mg twice a day.

	Notes		Start anning and		"Shil wit fil? Drielly the rule, Kolowe aspette.	Freeling better today.	"EQUIPATE MASE experyer pr. pr.	*Lacib at charak.
	Bhood Sugar		***				129 17 p.m.	
Jeetime	Blood Sugar	211	jų.	12,	2147.	S.	*788	252
Dinner	Blood Sugar	121		9 2	*	æ	125	, X84
Limit	Blood Sugar	384	6 62	22	129	葉		##
Breakfart	Blood Sugar	as.	211	類	##	*	RI	927
		noth	səqL	pays	swift	μ,	JPS	ung

Glucose Log Sheet for People Who Do Not Use Insulin Use this log sheet—or one like it that your health care provider may give you—to keep a record of your daily blood glucose levels.

Daily Log

Week Starting_

	Notes							
Other	Blood Sugar							
Bedtime	Blood Sugar							
	Blood Sugar							
Lumch	Sugar Sugar							
Dreakfast	Blood Sugar							
		иоуу	जिल्ह	₽₹M	enst!	Щ	₽S	ung

Glucose Log Sheet for People Who Use Insulin Use this log sheet—or one like it that your health care provider may give you—to keep a record of your daily blood glucose levels.

May 26, 2001

		***	ape, jein.			7#4E.	Į.			7	À			_146	i
1	Notes	"Worked and they	and-Oracle orașe,			Vine at Above	amerika, Kathari ngatin			1.04 Sant	Annie - Surg			"Sheeke	
Het.	Blood		W. C.			2	4//	0.							
Other	Dose			***************************************					3						
į	Blood Sugar	3	£	į	3	3	747	- 1	\$	ļ	E	į	7	\$	È
Bedtime	Dose		ø		4	•	4		*		4		P		*
Ter	poorg	ş	2//	1	8		ğ	1	?		8	3	*	1	į
Dinner	Dose	•		+		•	7	•		•		+		•	
Ę	Blood	+		B B		ş	5 8		ŧ	1	***	***	è	8	È.
Lunch	Dose	\$		4		*		2		7		3		7	
defant	Blood Sugar	#1		ħ		99				¥		*		į,	ŧ.
FIER	Dose	٠	R	*	R	•	R	8	R		P	4	Ą	6	R
9	luani ayt	A	#6#	Ay.	HUH	Ks.	HUH	68	HHH	P.	<i>₩#</i>	hy	11611	ks.	1100
not		W	521	Л	pą	М	SATI	ЦĽ	μ	ď	JB	5	un	5	

Glucose Log Sheet for People Who Use Insulin

Use this log sheet—or one like it that your health care provider may give you—to keep a record of your daily blood glucose levels.

Scotland State Week Starting_ Bedtine Blood Slood Dinner pooli Blood South Daily Log niluani eqq иоуу ævil SURLL Ш wig Pavi æς

Your Health Care Team

Primary Doctor or Health Care Provider

Name:
Telephone number:
Your questions:
Important points:

Eye Doctor (Ophthalmologist, Optometrist)

Name:
Telephone number:
Vour questions:
Your questions:
Important points:

Foot Doctor (Podiatrist)

Name:
Telephone number:
Variation and the same
Your questions:
Important points:

Dentist
Name:
Telephone number:
Your questions:
·
Important points:

Dietitian (Registered Dietitian, Nutritionist)

Name:
Telephone number:
Your questions:
Important points:

Diabetes Educator (Certified Diabetes Educator)

Name:
Telephone number:
Your questions:
Important points:

Counselor Name: Telephone number: ______ Your questions: _____ Important points:

Other
Name:
Telephone number:
Your questions:
Important points:

Glossary

A1C—A test that sums up how much glucose has been sticking to part of the hemoglobin during the past 3–4 months. Hemoglobin is a substance in the red blood cells that supplies oxygen to the cells of the body. the AIC goal for patients in general is an AIC goal of less than 7%. The AIC goal for the individual patient is an AIC as close to 6% as possible without a considerable amount of low blood glucose.

ACE inhibitor—A type of drug used to lower blood pressure. Studies indicate that it may also help prevent or slow the progression of kidney disease in people with diabetes. ACE is an acronym for angiotensin-converting enzyme.

autoimmune process—A process where the body's immune system attacks and destroys body tissue that it mistakes for foreign matter.

beta cells—Cells that make insulin. Beta cells are found in areas of the pancreas called the Islets of Langerhans.

bladder—A hollow organ that urine drains into from the kidneys. From the bladder, urine leaves the body.

blood glucose—The main sugar that the body makes from the food we eat. Glucose is carried through the bloodstream to provide energy to all of the body's living cells. The cells cannot use glucose without the help of insulin.

blood sugar—See blood glucose.

calluses—Thick, hardened areas of the skin, generally on the foot, caused by friction or pressure. Calluses can lead to other problems, including serious infection and even gangrene.

carbohydrate—One of three main groups of foods in the diet that provide calories and energy. (Protein and fat are the others.) Carbohydrates are mainly sugars (simple carbohydrates) and starches (complex carbohydrates, found in bread, pasta, beans) that the body breaks down into glucose.

cholesterol—A substance similar to fat that is found in the blood, muscles, liver, brain, and other body tissues. The body produces and needs some cholesterol. However, too much cholesterol can make fats stick to the walls of the arteries and cause a disease that decreases or stops circulation.

chronic kidney disease (CKD)—retains fluids and harmul wastes build up because the kidneys no longer work properly.

corns—A thickening of the skin of the feet or hands, usually caused by pressure against the skin.

dehydration—the loss of too much body fluid through frequent urinating, sweating, diarrhea, or vomiting.

diabetes—The short name for the disease called diabetes mellitus. Diabetes results when the body cannot use blood glucose as energy because of having too little insulin or being unable to use insulin. See also type 1 diabetes, type 2 diabetes, and gestational diabetes.

diabetes pills—Pills or capsules that are taken by mouth to help lower the blood glucose level. These pills may work for people whose bodies are still making insulin.

diabetic eye disease—A disease of the small blood vessels of the retina of the eye in people with diabetes. In this disease, the vessels swell and leak liquid into the retina, blurring the vision and sometimes leading to blindness.

diabetic ketoacidosis—High blood glucose with the presence of ketones in the urine and bloodstream, often caused by taking too little insulin or during illness.

diabetic kidney disease—Damage to the cells or blood vessels of the kidney.

diabetic nerve damage—Damage to the nerves of a person with diabetes. Nerve damage may affect the feet and hands, as well as major organs.

dialysis—A method for removing waste from the blood when the kidneys can no longer do the job.

dilated eye exam—Eye drops are placed in the eyes to widen the pupils to see the retina better. The eye doctor will look for changes in the retinain the back of the eyes.

diphtheria—An acute, contagious disease that causes fever and problems for the heart and nervous system.

EKG—A test that measures the heart's action. Also called an electrocardiogram.

flu—An infection caused by the "flu" (short for "influenza") virus. The flu is a contagious viral illness that strikes quickly and severely. Signs include high fever, chills, body aches, runny nose, sore throat, and headache.

gestational diabetes—A type of diabetes that can occur in pregnant women who have not been known to have diabetes before.

GFR—glumerular filration rate- A measure of the kidney's ability to filter and remove waste products. It is the best tst to measure kidney function and stage of kidney disease.

gingivitis—A swelling and soreness of the gums that, without treatment, can cause serious gum problems and disease.

glucagon—A hormone that raises the blood glucose level.

glucose—A sugar in our blood and a source of energy for our bodies.

heart attack—Damage to the heart muscle caused when the blood vessels supplying the muscle are blocked, such as when the blood vessels are clogged with fats (a condition sometimes called hardening of the arteries). **HDL** (or **high-density lipoprotein**)—A combined protein and fatlike substance. Low in cholesterol, it usually passes freely through the arteries. Sometimes called "good cholesterol."

high blood glucose—A condition that occurs in people with diabetes when their blood glucose levels are too high. Symptoms include having to urinate often, being very thirsty, and losing weight.

high blood pressure—A condition where the blood circulates through the arteries with too much force. High blood pressure tires the heart, harms the arteries, and increases the risk of heart attack, stroke, and kidney problems.

hormone—A chemical that special cells in the body release to help other cells work. For example, insulin is a hormone made in the pancreas to help the body use glucose as energy.

hyperglycemia—See high blood glucose.

hypertension—See high blood pressure.

hypoglycemia—See low blood glucose.

immunization—Sometimes called vaccination; a shot or injection that protects a person from getting an illness by making the person "immune" to it.

influenza—See flu.

inject—To force a liquid into the body with a needle and syringe.

insulin—A hormone that helps the body use blood glucose for energy. The beta cells of the pancreas make insulin. When people with diabetes can't make enough insulin, they may have to inject it from another source.

insulin-dependent diabetes—See type 1 diabetes.

ketones—Chemical substances that the body makes when it doesn't have enough insulin in the blood. When ketones build up in the body for a long time, serious illness or coma can result.

kidneys—Twin organs found in the lower part of the back. The kidneys purify the blood of all waste and harmful material. They also control the level of some helpful chemical substances in the blood.

laser surgery—Surgery that uses a strong ray of special light, called a laser, to treat damaged parts of the body. Laser surgery can help treat some diabetic eye diseases.

low blood glucose—A condition that occurs in people with diabetes when their blood glucose levels are too low. Symptoms include feeling anxious or confused, feeling numb in the arms and hands, and shaking or feeling dizzy.

LDL (or **low-density lipoprotein)**—A combined protein and fatlike substance. Rich in cholesterol, it tends to stick to the walls in the arteries. Sometimes called "bad cholesterol."

meal plan—A guide to help people get the proper amount of calories, carbohydrates, proteins, and fats in their diet. See also food exchanges.

microalbumin—A protein found in blood plasma and urine. The presence of microalbumin in the urine can be a sign of kidney disease.

nephropathy—See diabetic kidney disease.

neuropathy—See diabetic nerve damage.

non-insulin-dependent diabetes—See type 2 diabetes.

pancreas—An organ in the body that makes insulin so that the body can use glucose for energy. The pancreas also makes enzymes that help the body digest food.

periodontitis—A gum disease in which the gums shrink away from the teeth. Without treatment, it can lead to tooth loss.

plaque—A film of mucus that traps bacteria on the surface of the teeth. Plaque can be removed with daily brushing and flossing of teeth.

pumice stone—A special foot care tool used to gently file calluses as instructed by your health care team.

retinopathy—See diabetic eye disease.

risk factors—Traits that make it more likely that a person will get an illness. For example, a risk factor for getting type 2 diabetes is having a family history of diabetes.

self-monitoring blood glucose—A way for people with diabetes to find out how much glucose is in their blood. A drop of blood from the fingertip is placed on a special coated strip of paper that "reads" (often through an electronic meter) the amount of glucose in the blood.

stroke—Damage to a part of the brain that happens when the blood vessels supplying that part are blocked, such as when the blood vessels are clogged with fats (a condition sometimes called hardening of the arteries).

support group—A group of people who share a similar problem or concern. The people in the group help one another by sharing experiences, knowledge, and information.

type 1 diabetes—A condition in which the pancreas makes so little insulin that the body can't use blood glucose as energy. People with type 1 diabetes ned to take insulin every day.

type 2 diabetes—A condition in which the body either makes too little insulin or can't use the insulin it makes to use blood glucose as energy. All people with diabetes need to eat healthy foods stay at a healthy weight and be active everyday. People with type 2 often need to diabetes have to take diabetes pills or insulin. type 2 diabetes is the most common from of diabetes.

ulcer —A break or deep sore in the skin. Germs can enter an ulcer and may be hard to heal.

urea—One of the chief waste products of the body. When the body breaks down food, it uses what it needs and throws the rest away as waste. The kidneys flush the waste from the body in the form of urea, which is in the urine.

vaccination—A shot given to protect against a disease.

vagina—A canal in females from the external genitalia (vulva) to the cervix of the uterus.

vitrectomy—An operation to remove the blood that sometimes collects at the back of the eyes when a person has eye disease.

yeast infection —A vaginal infection that is usually caused by a fungus. Women who have this infection may feel itching, burning when urinating, and pain, and some women have a vaginal discharge. Yeast infections occur more frequently in women with diabetes.

Resources

The following is a list of organizations that can provide information on diabetes. Ask your health care team to help you find other resources for information or support.

American Association of Diabetes Educators 100 West Monroe, Suite 400 Chicago, Illinois 60603-1901 800-832-6874 800-338-3633 (for names of diabetes educators) http://www.diabeteseducator.org

American Diabetes Association
1701 N. Beauregard Street
Alexandria, Virginia 22311
703-549-1500
800-ADA-ORDER to order publicationstoll free
800-342-2388 (800-DIABETES) for diabetes
information
http://www.diabetes.org

American Dietetic Association
216 West Jackson Boulevard, Suite 800
Chicago, Illinois 60606-6995
800-745-0775
800-366-1655 (Consumer Nutrition Hotline,
Spanish speaker available)
http://www.eatright.org

American Heart Association National Center 7272 Greenville Avenue Dallas, Texas 75231 214-373-6300 http://www.americanheart.org

Centers for Disease Control and Prevention (CDC)
Division of Diabetes Translation
1-800-CDC-INFO (232-4636)
E-mail: cdcinfo@cdc.gov
http://www.cdc.gov/diabetes

Department of Veterans Affairs

Veterans Health Administration Diabetes Program http://www.va.gov/diabetes
Veterans Health Administration
810 Vermont Avenue, N.W.
Washington, DC 20420
VA Health Benefits
Toll-free: 877-222-8387
https://iris.va.gov/phonenbrs.asp

The Foundation of the American Academy of Ophthalmology Diabetes Project P.O. Box 429098 San Francisco, California 94142-9098 800-222-EYES (3937) http://www.aao.org/aaoweb1/foundation/301.cfm

Indian Health Service Diabetes Program 5300 Homestead Road, N.E. Albuquerque, New Mexico 87110 505-248-4182 http://www.ihs.gov/medicalprograms/diabetes

Juvenile Diabetes Research Foundation International The Diabetes Research Foundation 120 Wall Street, 19th Floor New York, New York 10005-4001 800-JDF-CUREor 800-223-1138 212-785-9595 (fax) E-mail: info@jdrf.org http://www.jdf.org

National Diabetes Education Program

• Program information:

Mail requests to NDEP, c/o CDC Diabetes Program Public Inquiries

P.O. Box 8728

Silver Spring, Maryland 20910

877-CDC-DIAB

800-438-5383

E-mail: diabetes@cdc.gov

http://www.cdc.gov/diabetes/ndep/index.htm

· Campaign materials and publications:

Mail requests to NDEP, c/o National Diabetes Information Clearinghouse

1 Information Way

Bethesda, Maryland 20892-3560

800-438-5383

E-mail: berryt@extra.niddk.nih.gov

http://www.ndep.nih.gov

National Diabetes Information Clearinghouse

1 Information Way

Bethesda, Maryland 20892-3560

800-860-8747

301-654-3327

Fax 301-907-8906

E-mail: ndic@aerie.com

http://www.diabetes.niddk.nih.gov/index.htm

National Eye Institute

National Eye Health Education Program

Diabetic Eye Disease Public Education Program

2020 Vision Place

Bethesda, Maryland 20892-3655

800-869-2020 (to order materials)

Others:	 	 	

Others: _			