

Recall the unit circle $x^2 + y^2 = 1^2$

$$\sin \theta = y$$

$$\cos \theta = x$$

$$\tan \theta = \frac{y}{x}, x \neq 0$$

$$\csc \theta = \frac{1}{y}, y \neq 0$$

$$\sec \theta = \frac{1}{x}, x \neq 0$$

$$\cot \theta = \frac{x}{y}, y \neq 0$$

The following can be inferred...

$$\sin \theta = \frac{1}{\csc \theta}$$

$$\csc \theta = \frac{1}{\sin \theta}$$

$$\cos \theta = \frac{1}{\sec \theta}$$

$$\sec \theta = \frac{1}{\cos \theta}$$

$$\tan \theta = \frac{\sin \theta}{\cos \theta} = \frac{1}{\cot \theta}$$

$$\cot \theta = \frac{\cos \theta}{\sin \theta} = \frac{1}{\tan \theta}$$

These are our "basic trig identities".

$\rightarrow x^2 + y^2 = 1$ ← definition of unit circle.

Recall $x = \cos \theta$, $y = \sin \theta$

$$(\cos \theta)^2 + (\sin \theta)^2 = 1$$

$$\cos^2 \theta + \sin^2 \theta = 1$$

"Pythagorean Identity"

$$\hookrightarrow \cos^2 \theta = 1 - \sin^2 \theta$$

$$\hookrightarrow \sin^2 \theta = 1 - \cos^2 \theta$$

Note: e.g.

$$\cos^2 \theta = (\cos \theta)^2$$

$$\frac{\tan^6 \theta = (\tan \theta)^6}{\text{However}}$$

$$\sin^{-1} \theta \neq (\sin \theta)^{-1} \\ \neq \frac{1}{\sin \theta}$$

$$\sin^{-1} \theta = \arcsin \theta$$

$$\frac{\cos^2 \theta}{\cos^2 \theta} + \frac{\sin^2 \theta}{\cos^2 \theta} = \frac{1}{\cos^2 \theta}$$

$$1 + \left(\frac{\sin \theta}{\cos \theta}\right)^2 = \left(\frac{1}{\cos \theta}\right)^2$$

$$1 + (\tan \theta)^2 = (\sec \theta)^2$$

$$1 + \tan^2 \theta = \sec^2 \theta$$

$$\hookrightarrow \tan^2 \theta = \sec^2 \theta - 1$$

$$\hookrightarrow 1 = \sec^2 \theta - \tan^2 \theta$$

$$\frac{\cos^2 \theta}{\sin^2 \theta} + \frac{\sin^2 \theta}{\sin^2 \theta} = \frac{1}{\sin^2 \theta}$$

$$\left(\frac{\cos \theta}{\sin \theta}\right)^2 + 1 = \left(\frac{1}{\sin \theta}\right)^2$$

$$(\cot \theta)^2 + 1 = (\csc \theta)^2$$

$$\cot^2 \theta + 1 = \csc^2 \theta$$

$$\hookrightarrow \cot^2 \theta = \csc^2 \theta - 1$$

$$\hookrightarrow 1 = \csc^2 \theta - \cot^2 \theta$$



Identity: if given two functions $f(x)$ and $g(x)$

$f(x) = g(x)$ for all values of x in both functions' domains.

Consider the system of 2 linear equations

$$y = 2x - 1$$

$$y = 2(x - 3) + 5$$



Both certainly look equal. Prove it.

$$2x - 1 = 2(x - 3) + 5$$

$$2x - 1 = 2x - 6 + 5$$

$$2x - 1 = 2x - 1 \quad \square \text{ or QED}$$

Notice, we DO NOT MOVE ANYTHING OVER THE EQUAL SIGN.

Our objective for this topic is to prove that these identities are true. We will be using various algebraic manipulations and identity substitutions in order to do so.

I repeat, DO NOT MOVE ANYTHING OVER THE EQUAL SIGN.

Verify that these equations are identities.

$$\sin \theta \cot \theta = \cos \theta$$

$$\sin \theta \left(\frac{\cos \theta}{\sin \theta} \right) = \cos \theta$$

$$\frac{\sin \theta \cos \theta}{\sin \theta} = \cos \theta$$

$$\cos \theta = \cos \theta \quad \square$$

* Consider yourselves
to be
"plastic surgeons"
Your job is to
"make ugly into pretty"

$$\cos \theta (\sec \theta - \cos \theta) = \sin^2 \theta$$

$$\cos \theta \cdot \sec \theta - \cos \theta \cdot \cos \theta = \sin^2 \theta$$

$$\cos \theta \left(\frac{1}{\cos \theta} \right) - (\cos^2 \theta) = \sin^2 \theta$$

$$1 - \cos^2 \theta = \sin^2 \theta$$

$$(\cos^2 \theta + \sin^2 \theta) - \cos^2 \theta = \sin^2 \theta$$

$$\sin^2 \theta = \sin^2 \theta$$

$$1 - (1 - \sin^2 \theta) = \sin^2 \theta$$

also viable

QED

$$\tan \theta = \frac{\sin \theta \cos \theta + \sin \theta}{\cos \theta + \cos^2 \theta}$$

$$\tan \theta = \frac{\sin \theta (\cos \theta + 1)}{\cos \theta (1 + \cos \theta)}$$

$$\tan \theta = \frac{\sin \theta}{\cos \theta}$$

$$\tan \theta = \tan \theta \quad \square$$

$$\underbrace{(1 + \sin \theta)}_{=} \underbrace{(1 - \sin \theta)}_{=} = \cos^2 \theta$$

$a^2 - b^2 = (a+b)(a-b)$

$$1^2 - \sin^2 \theta = \cos^2 \theta$$

$$1 - \sin^2 \theta = \cos^2 \theta$$

$$\cos^2 \theta = \cos^2 \theta$$

QED

$$\frac{(\sin\theta + \cos\theta)^2}{\cos\theta} = \sec\theta + 2\sin\theta$$

$$(a+b)^2 = a^2 + 2ab + b^2$$

$$\frac{(\sin\theta + \cos\theta)(\sin\theta + \cos\theta)}{\cos\theta} = \sec\theta + 2\sin\theta$$

$$\frac{\sin^2\theta + 2\sin\theta\cos\theta + \cos^2\theta}{\cos\theta} = \sec\theta + 2\sin\theta$$

$$\frac{(\sin^2\theta + \cos^2\theta) + 2\sin\theta\cos\theta}{\cos\theta} = \sec\theta + 2\sin\theta$$

$$\frac{1 + 2\sin\theta\cos\theta}{\cos\theta} = \sec\theta + 2\sin\theta$$

$$\frac{1}{\cos\theta} + \frac{2\sin\theta\cos\theta}{\cos\theta} = \sec\theta + 2\sin\theta$$

$$\sec\theta + 2\sin\theta = \sec\theta + 2\sin\theta$$

$$\frac{\cos^2 \theta}{\sin \theta} + \sin \theta = \csc \theta$$

$$\frac{\cos^2 \theta}{\sin \theta} + \sin \theta \left(\frac{\sin \theta}{\sin \theta} \right) = \frac{1}{\sin \theta}$$

$$\frac{\cos^2 \theta}{\sin \theta} + \frac{\sin^2 \theta}{\sin \theta} = \frac{1}{\sin \theta}$$

$$\frac{\cos^2 \theta + \sin^2 \theta}{\sin \theta} = \frac{1}{\sin \theta}$$

$$\frac{1}{\sin \theta} = \frac{1}{\sin \theta}$$

QED

$$\sin^2 \theta \tan^2 \theta = \tan^2 \theta - \sin^2 \theta$$

$$\frac{\cos(t)}{1 + \sec(t)} = \frac{1 - \cos(t)}{\tan^2(t)}$$

$$\left(\frac{\cos(t)}{1 + \sec(t)} \right) \cdot \left(\frac{1 - \sec(t)}{1 - \sec(t)} \right) = \frac{1 - \cos(t)}{\tan^2(t)}$$

conjugate

$$\frac{\cos(t) - \cos(t) \sec(t)}{1 - \sec^2(t)} = \frac{1 - \cos(t)}{\tan^2(t)}$$

$$\frac{\cos(t) - \cos(t) \left(\frac{1}{\cos(t)} \right)}{1 - \sec^2(t)} = \frac{1 - \cos(t)}{\tan^2(t)}$$

$$\frac{\cos(t) - 1}{1 - \sec^2(t)} = \frac{1 - \cos(t)}{\sec^2(t) - 1}$$

$$\frac{-1(-\cos(t) + 1)}{-1(-1 + \sec^2(t))} = \frac{1 - \cos(t)}{\sec^2(t) - 1}$$

$$\frac{1 - \cos(t)}{\sec^2(t) - 1} = \frac{1 - \cos(t)}{\sec^2(t) - 1} \quad \square$$

$$\frac{\sec(x)}{\sin(x)} - \frac{\sin(x)}{\sec(x)} = \frac{\tan^2(x) + \cos^2(x)}{\tan(x)}$$

Modullell.

$$1. \quad 1 + \cos A = \frac{\sin^2 A}{1 - \cos A} \cdot \left(\frac{1 + \cos A}{1 + \cos A} \right)$$

$$1 + \cos A = \frac{\sin^2 A (1 + \cos A)}{1 - \cos^2 A}$$

$$1 + \cos A = \frac{\sin^2 A (1 + \cos A)}{(\sin^2 A)}$$

$$1 + \cos A = 1 + \cos A \quad \square$$

$$4. \quad \sin^2(\alpha) = \frac{1 - \cos^4(\alpha)}{1 + \cos^2(\alpha)} \quad \rightarrow \quad 1 - \cos^4(\alpha) = (1)^2 - (\cos^2(\alpha))^2$$

$$\sin^2(\alpha) = \frac{(1 + \cos^2(\alpha)) (1 - \cos^2(\alpha))}{(1 + \cos^2(\alpha))}$$

$$\sin^2(\alpha) = 1 - \cos^2(\alpha)$$

$$\sin^2(\alpha) = \sin^2(\alpha) \quad \square$$

7.

$$2 \csc^2(t) = \frac{1}{1 - \cos(t)} + \frac{1}{1 + \cos(t)}$$

$$2 \csc^2(t) = \frac{1}{1 - \cos(t)} \overset{\text{LCD}}{\left(\frac{1 + \cos(t)}{1 + \cos(t)} \right)} + \frac{1}{1 + \cos(t)} \overset{\text{LCD}}{\left(\frac{1 - \cos(t)}{1 - \cos(t)} \right)}$$

$$2 \csc^2(t) = \frac{1 + \cos(t) + 1 - \cos(t)}{(1 - \cos(t))(1 + \cos(t))}$$

$$2 \csc^2(t) = \frac{2}{1 - \cos^2(t)}$$

$$2 \csc^2(t) = \frac{2}{\sin^2(t)}$$

$$2 \csc^2(t) = 2 \left(\frac{1}{\sin^2(t)} \right)$$

$$2 \csc^2(t) = 2 \csc^2(t) \quad \text{Q.E.D.}$$