

Trigonometric functions reviewed

Lesson #17

MAT 1375 Precalculus


New York City College of Technology CUNY



©Thomas Tradler, Kate Poirier


Angles in degree and radian - review

Degree and radian




Angles in degree and radian - review

Degree and radian




Angles in degree and radian - review

Degree and radian




Angles in degree and radian - review

Degree and radian




Angles in degree and radian - review

Degree and radian




Angles in degree and radian - review

Degree and radian



Angles in degree and radian - review

Degree and radian



Conversion between degree and radian


$$180^\circ = \pi$$

or

$$1^\circ = \frac{\pi}{180}$$

Angles in degree and radian - review

Degree and radian



Convert to degree.

1 $\frac{\pi}{3} =$

2 $\frac{11\pi}{6} =$

3 $\frac{7\pi}{4} =$

4 $\frac{5\pi}{4} =$

5 $\frac{3\pi}{2} =$

Conversion between degree and radian


$$180^\circ = \pi$$

or

$$1^\circ = \frac{\pi}{180}$$

Angles in degree and radian - review

Degree and radian



Convert to degree.

- 1 $\frac{\pi}{3} = \frac{180^\circ}{3} = 60^\circ$
- 2 $\frac{11\pi}{6} =$
- 3 $\frac{7\pi}{4} =$
- 4 $\frac{5\pi}{4} =$
- 5 $\frac{3\pi}{2} =$

Conversion between degree and radian


$$180^\circ = \pi$$

or

$$1^\circ = \frac{\pi}{180}$$

Angles in degree and radian - review

Degree and radian



Convert to degree.

① $\frac{\pi}{3} = \frac{180^\circ}{3} = 60^\circ$

② $\frac{11\pi}{6} = \frac{11 \cdot 180^\circ}{6} = 11 \cdot 30^\circ = 330^\circ$

③ $\frac{7\pi}{4} =$

④ $\frac{5\pi}{4} =$

⑤ $\frac{3\pi}{2} =$

Conversion between degree and radian


$$180^\circ = \pi$$

or

$$1^\circ = \frac{\pi}{180}$$

Angles in degree and radian - review

Degree and radian



Convert to degree.

$$① \quad \frac{\pi}{3} = \frac{180^\circ}{3} = 60^\circ$$

$$② \quad \frac{11\pi}{6} = \frac{11 \cdot 180^\circ}{6} = 11 \cdot 30^\circ = 330^\circ$$

$$③ \quad \frac{7\pi}{4} = \frac{7 \cdot 180^\circ}{4} = 7 \cdot 45^\circ = 315^\circ$$

$$④ \quad \frac{5\pi}{4} = \frac{5 \cdot 180^\circ}{4} = \frac{900^\circ}{4} = 225^\circ$$

$$⑤ \quad \frac{3\pi}{2} = \frac{3 \cdot 180^\circ}{2} = 3 \cdot 90^\circ = 270^\circ$$

Conversion between degree and radian


$$180^\circ = \pi$$

or

$$1^\circ = \frac{\pi}{180}$$

Angles in degree and radian - review

Degree and radian



Conversion between degree and radian

$$180^\circ = \pi$$

or

$$1^\circ = \frac{\pi}{180}$$

Convert to degree.

① $\frac{\pi}{3} = \frac{180^\circ}{3} = 60^\circ$

② $\frac{11\pi}{6} = \frac{11 \cdot 180^\circ}{6} = 11 \cdot 30^\circ = 330^\circ$

③ $\frac{7\pi}{4} = \frac{7 \cdot 180^\circ}{4} = 7 \cdot 45^\circ = 315^\circ$

④ $\frac{5\pi}{4} = \frac{5 \cdot 180^\circ}{4} = \frac{900^\circ}{4} = 225^\circ$

⑤ $\frac{3\pi}{2} = \frac{3 \cdot 180^\circ}{2} = 3 \cdot 90^\circ = 270^\circ$

Convert to radian.

① $120^\circ =$

② $300^\circ =$

③ $210^\circ =$


④ $150^\circ =$

⑤ $135^\circ =$

⑥ $240^\circ =$

Angles in degree and radian - review

Degree and radian



Conversion between degree and radian

$$180^\circ = \pi$$

or

$$1^\circ = \frac{\pi}{180}$$

Convert to degree.

$$① \quad \frac{\pi}{3} = \frac{180^\circ}{3} = 60^\circ$$

$$② \quad \frac{11\pi}{6} = \frac{11 \cdot 180^\circ}{6} = 11 \cdot 30^\circ = 330^\circ$$

$$③ \quad \frac{7\pi}{4} = \frac{7 \cdot 180^\circ}{4} = 7 \cdot 45^\circ = 315^\circ$$

$$④ \quad \frac{5\pi}{4} = \frac{5 \cdot 180^\circ}{4} = \frac{900^\circ}{4} = 225^\circ$$

$$⑤ \quad \frac{3\pi}{2} = \frac{3 \cdot 180^\circ}{2} = 3 \cdot 90^\circ = 270^\circ$$

Convert to radian.

$$① \quad 120^\circ = 120 \cdot \frac{\pi}{180} = \frac{2\pi}{3}$$

$$② \quad 300^\circ =$$

$$③ \quad 210^\circ =$$


$$④ \quad 150^\circ =$$

$$⑤ \quad 135^\circ =$$

$$⑥ \quad 240^\circ =$$

Angles in degree and radian - review

Degree and radian



Conversion between degree and radian

$$180^\circ = \pi$$

or

$$1^\circ = \frac{\pi}{180}$$

Convert to degree.

$$① \quad \frac{\pi}{3} = \frac{180^\circ}{3} = 60^\circ$$

$$② \quad \frac{11\pi}{6} = \frac{11 \cdot 180^\circ}{6} = 11 \cdot 30^\circ = 330^\circ$$

$$③ \quad \frac{7\pi}{4} = \frac{7 \cdot 180^\circ}{4} = 7 \cdot 45^\circ = 315^\circ$$

$$④ \quad \frac{5\pi}{4} = \frac{5 \cdot 180^\circ}{4} = \frac{900^\circ}{4} = 225^\circ$$

$$⑤ \quad \frac{3\pi}{2} = \frac{3 \cdot 180^\circ}{2} = 3 \cdot 90^\circ = 270^\circ$$

Convert to radian.

$$① \quad 120^\circ = 120 \cdot \frac{\pi}{180} = \frac{2\pi}{3}$$

$$② \quad 300^\circ = 300 \cdot \frac{\pi}{180} = \frac{5\pi}{3}$$


$$③ \quad 210^\circ = 210 \cdot \frac{\pi}{180} = \frac{7\pi}{6}$$

$$④ \quad 150^\circ = 150 \cdot \frac{\pi}{180} = \frac{5\pi}{6}$$


$$⑤ \quad 135^\circ = 135 \cdot \frac{\pi}{180} = \frac{3\pi}{4}$$

$$⑥ \quad 240^\circ = 240 \cdot \frac{\pi}{180} = \frac{4\pi}{3}$$

Trigonometric functions via unit circle - the idea




Trigonometric functions via unit circle - the idea



sin:

The silver sphere figure measures the vertical position of the passenger car from the center.

Trigonometric functions via unit circle - the idea




sin:

The silver sphere figure measures the vertical position of the passenger car from the center.

cos:

The copper cone figure measures the horizontal position of the passenger car from the center.

Trigonometric functions via unit circle - the idea



sin:

The silver sphere figure measures the vertical position of the passenger car from the center.

cos:

The copper cone figure measures the horizontal position of the passenger car from the center.


* Note:

The wonder circle has radius 1.

Trigonometric functions via reference angles - review

Definition (Trigonometric functions)

For a point (x, y) on the unit circle, which is on the terminal side of θ :



$$\sin(\theta) = y$$


$$\cos(\theta) = x$$

$$\tan(\theta) = \frac{\sin(\theta)}{\cos(\theta)} = \frac{y}{x}$$

Trigonometric functions via reference angles - review

Definition (Trigonometric functions)

For a point (x, y) on the unit circle, which is on the terminal side of θ :




$$\sin(\theta) = y$$

$$\cos(\theta) = x$$

$$\tan(\theta) = \frac{\sin(\theta)}{\cos(\theta)} = \frac{y}{x}$$

Reference angle

The **reference angle** is the smallest angle that the terminal side forms with the x -axis.




reference angle:

$$\frac{\pi}{6}$$

Trigonometric functions via reference angles - review

Definition (Trigonometric functions)

For a point (x, y) on the unit circle, which is on the terminal side of θ :




$$\sin(\theta) = y$$

$$\cos(\theta) = x$$

$$\tan(\theta) = \frac{\sin(\theta)}{\cos(\theta)} = \frac{y}{x}$$

Reference angle

The **reference angle** is the smallest angle that the terminal side forms with the x -axis.



reference angle:


$$\frac{\pi}{6}$$

$$\frac{\pi}{4}$$

Trigonometric functions via reference angles - review

Definition (Trigonometric functions)

For a point (x, y) on the unit circle, which is on the terminal side of θ :




$$\sin(\theta) = y$$

$$\cos(\theta) = x$$

$$\tan(\theta) = \frac{\sin(\theta)}{\cos(\theta)} = \frac{y}{x}$$

Reference angle

The **reference angle** is the smallest angle that the terminal side forms with the x -axis.



reference angle:

$$\frac{\pi}{6}$$


$$\frac{\pi}{4}$$

$$\frac{\pi}{3}$$

Trigonometric functions via reference angles - review

Definition (Trigonometric functions)

For a point (x, y) on the unit circle, which is on the terminal side of θ :




$$\sin(\theta) = y$$

$$\cos(\theta) = x$$

$$\tan(\theta) = \frac{\sin(\theta)}{\cos(\theta)} = \frac{y}{x}$$


Reference angle

The **reference angle** is the smallest angle that the terminal side forms with the x -axis.



Coordinates on the unit circle

Coordinates coincide with the reference angle up to “ \pm ” sign.




Thus: \sin , \cos , \tan of an angle coincide with those of the reference angle except for “ \pm ” sign.

Trigonometric functions via reference angles - review

Definition (Trigonometric functions)

For a point (x, y) on the unit circle, which is on the terminal side of θ :




$$\sin(\theta) = y$$

$$\cos(\theta) = x$$

$$\tan(\theta) = \frac{\sin(\theta)}{\cos(\theta)} = \frac{y}{x}$$

Reference angle

The **reference angle** is the smallest angle that the terminal side forms with the x -axis.



reference angle:


$$\frac{\pi}{6}$$

$$\frac{\pi}{4}$$

$$\frac{\pi}{3}$$

Coordinates on the unit circle

Coordinates coincide with the reference angle up to “ \pm ” sign.




Thus: \sin , \cos , \tan of an angle coincide with those of the reference angle except for “ \pm ” sign.

Trigonometric functions via reference angles - review

Definition (Trigonometric functions)

For a point (x, y) on the unit circle, which is on the terminal side of θ :




$$\sin(\theta) = y$$

$$\cos(\theta) = x$$

$$\tan(\theta) = \frac{\sin(\theta)}{\cos(\theta)} = \frac{y}{x}$$

Reference angle

The **reference angle** is the smallest angle that the terminal side forms with the x-axis.



reference angle:


$$\frac{\pi}{6}$$

$$\frac{\pi}{4}$$

$$\frac{\pi}{3}$$

Coordinates on the unit circle

Coordinates coincide with the reference angle up to “ \pm ” sign.




Thus: \sin , \cos , \tan of an angle coincide with those of the reference angle except for “ \pm ” sign.

Trigonometric functions via reference angles - review

Definition (Trigonometric functions)

For a point (x, y) on the unit circle, which is on the terminal side of θ :




$$\sin(\theta) = y$$

$$\cos(\theta) = x$$

$$\tan(\theta) = \frac{\sin(\theta)}{\cos(\theta)} = \frac{y}{x}$$

Reference angle

The **reference angle** is the smallest angle that the terminal side forms with the x -axis.



reference angle:


$$\frac{\pi}{6}$$

$$\frac{\pi}{4}$$

$$\frac{\pi}{3}$$

Coordinates on the unit circle

Coordinates coincide with the reference angle up to “ \pm ” sign.



Thus: \sin , \cos , \tan of an angle coincide with those of the reference angle except for “ \pm ” sign.

Using the calculator

$$\frac{1}{2} = 0.5$$

$$1 = 1$$

$$\frac{\sqrt{2}}{2} \approx 0.707$$


$$\sqrt{3} \approx 1.732$$

$$\frac{\sqrt{3}}{2} \approx 0.866$$

$$\frac{\sqrt{3}}{3} \approx 0.577$$

Trigonometric functions via reference angles - exercises

Trigonometric functions via reference angle



$$\begin{aligned} \frac{1}{2} &= 0.5 & 1 &= 1 \\ \frac{\sqrt{2}}{2} &\approx 0.707 & \sqrt{3} &\approx 1.732 \\ \frac{\sqrt{3}}{2} &\approx 0.866 & \frac{\sqrt{3}}{3} &\approx 0.577 \end{aligned}$$

1 Find $\sin(\frac{5\pi}{3})$.

2 Find $\cos(\frac{5\pi}{4})$.


4 Find $\sin(135^\circ)$.

3 Find $\sin(-\frac{11\pi}{6})$.

5 Find $\cos(210^\circ)$.

Trigonometric functions via reference angles - exercises

Trigonometric functions via reference angle



$$\frac{1}{2} = 0.5 \quad 1 = 1$$
$$\frac{\sqrt{2}}{2} \approx 0.707 \quad \sqrt{3} \approx 1.732$$
$$\frac{\sqrt{3}}{2} \approx 0.866 \quad \frac{\sqrt{3}}{3} \approx 0.577$$

1 Find $\sin(\frac{5\pi}{3})$.

Using calculator:

$$\sin(\frac{5\pi}{3}) \approx -0.866$$

2 Find $\cos(\frac{5\pi}{4})$.


4 Find $\sin(135^\circ)$.

3 Find $\sin(-\frac{11\pi}{6})$.

5 Find $\cos(210^\circ)$.

Trigonometric functions via reference angles - exercises

Trigonometric functions via reference angle



$$\cos \theta = x$$

$$\sin \theta = y$$

$$\tan \theta = \frac{y}{x}$$

$$\frac{1}{2} = 0.5$$

$$1 = 1$$

$$\frac{\sqrt{2}}{2} \approx 0.707$$

$$\sqrt{3} \approx 1.732$$

$$\frac{\sqrt{3}}{2} \approx 0.866$$

$$\frac{\sqrt{3}}{3} \approx 0.577$$

1 Find $\sin(\frac{5\pi}{3})$.

Using calculator:

$$\sin(\frac{5\pi}{3}) \approx -0.866$$

$$\Rightarrow \sin(\frac{5\pi}{3}) = -\frac{\sqrt{3}}{2}$$

2 Find $\cos(\frac{5\pi}{4})$.


4 Find $\sin(135^\circ)$.

3 Find $\sin(-\frac{11\pi}{6})$.

5 Find $\cos(210^\circ)$.

Trigonometric functions via reference angles - exercises

Trigonometric functions via reference angle



$$\frac{1}{2} = 0.5 \quad 1 = 1$$
$$\frac{\sqrt{2}}{2} \approx 0.707 \quad \sqrt{3} \approx 1.732$$
$$\frac{\sqrt{3}}{2} \approx 0.866 \quad \frac{\sqrt{3}}{3} \approx 0.577$$

1 Find $\sin(\frac{5\pi}{3})$.

Using calculator:

$$\sin(\frac{5\pi}{3}) \approx -0.866$$

$$\Rightarrow \sin(\frac{5\pi}{3}) = -\frac{\sqrt{3}}{2}$$

By hand: Reference angle: $\frac{\pi}{3}$

$$\sin(\frac{5\pi}{3}) = \pm \sin(\frac{\pi}{3}) = \pm \frac{\sqrt{3}}{2}$$

Quadrant of $\frac{5\pi}{3} = 300^\circ$: IV

2 Find $\cos(\frac{5\pi}{4})$.


4 Find $\sin(135^\circ)$.

3 Find $\sin(-\frac{11\pi}{6})$.

5 Find $\cos(210^\circ)$.

Trigonometric functions via reference angles - exercises

Trigonometric functions via reference angle



$$\frac{1}{2} = 0.5$$

$$1 = 1$$

$$\frac{\sqrt{2}}{2} \approx 0.707$$

$$\sqrt{3} \approx 1.732$$

$$\frac{\sqrt{3}}{2} \approx 0.866$$

$$\frac{\sqrt{3}}{3} \approx 0.577$$

1 Find $\sin(\frac{5\pi}{3})$.

Using calculator:

$$\sin(\frac{5\pi}{3}) \approx -0.866$$

$$\Rightarrow \sin(\frac{5\pi}{3}) = -\frac{\sqrt{3}}{2}$$

By hand: Reference angle: $\frac{\pi}{3}$

$$\sin(\frac{5\pi}{3}) = \pm \sin(\frac{\pi}{3}) = \pm \frac{\sqrt{3}}{2}$$

Quadrant of $\frac{5\pi}{3} = 300^\circ$: IV

2 Find $\cos(\frac{5\pi}{4})$.

Calculator:

$$\cos(\frac{5\pi}{4}) \approx -0.707$$


4 Find $\sin(135^\circ)$.

3 Find $\sin(-\frac{11\pi}{6})$.

5 Find $\cos(210^\circ)$.

Trigonometric functions via reference angles - exercises

Trigonometric functions via reference angle



$$\frac{1}{2} = 0.5$$

$$1 = 1$$

$$\frac{\sqrt{2}}{2} \approx 0.707$$

$$\sqrt{3} \approx 1.732$$

$$\frac{\sqrt{3}}{2} \approx 0.866$$

$$\frac{\sqrt{3}}{3} \approx 0.577$$

1 Find $\sin(\frac{5\pi}{3})$.

Using calculator:

$$\sin(\frac{5\pi}{3}) \approx -0.866$$

$$\Rightarrow \sin(\frac{5\pi}{3}) = -\frac{\sqrt{3}}{2}$$

By hand: Reference angle: $\frac{\pi}{3}$

$$\sin(\frac{5\pi}{3}) = \pm \sin(\frac{\pi}{3}) = \pm \frac{\sqrt{3}}{2}$$

Quadrant of $\frac{5\pi}{3} = 300^\circ$: IV

2 Find $\cos(\frac{5\pi}{4})$.

Calculator:

$$\cos(\frac{5\pi}{4}) \approx -0.707$$

$$\Rightarrow \cos(\frac{5\pi}{4}) = -\frac{\sqrt{2}}{2}$$


4 Find $\sin(135^\circ)$.

3 Find $\sin(-\frac{11\pi}{6})$.

5 Find $\cos(210^\circ)$.

Trigonometric functions via reference angles - exercises

Trigonometric functions via reference angle



$$\frac{1}{2} = 0.5$$

$$1 = 1$$

$$\frac{\sqrt{2}}{2} \approx 0.707$$

$$\sqrt{3} \approx 1.732$$

$$\frac{\sqrt{3}}{2} \approx 0.866$$

$$\frac{\sqrt{3}}{3} \approx 0.577$$

1 Find $\sin(\frac{5\pi}{3})$.

Using calculator:

$$\sin(\frac{5\pi}{3}) \approx -0.866$$

$$\Rightarrow \sin(\frac{5\pi}{3}) = -\frac{\sqrt{3}}{2}$$

By hand: Reference angle: $\frac{\pi}{3}$

$$\sin(\frac{5\pi}{3}) = \pm \sin(\frac{\pi}{3}) = \pm \frac{\sqrt{3}}{2}$$

Quadrant of $\frac{5\pi}{3} = 300^\circ$: IV

2 Find $\cos(\frac{5\pi}{4})$.

Calculator:

$$\cos(\frac{5\pi}{4}) \approx -0.707$$

$$\Rightarrow \cos(\frac{5\pi}{4}) = -\frac{\sqrt{2}}{2}$$

Reference angle: $\frac{\pi}{4}$

$$\cos(\frac{5\pi}{4}) = \pm \cos(\frac{\pi}{4}) = \pm \frac{\sqrt{2}}{2}$$

Quadrant of $\frac{5\pi}{4} = 225^\circ$: III


4 Find $\sin(135^\circ)$.

3 Find $\sin(-\frac{11\pi}{6})$.

5 Find $\cos(210^\circ)$.

Trigonometric functions via reference angles - exercises

Trigonometric functions via reference angle



$$\frac{1}{2} = 0.5$$

$$1 = 1$$

$$\frac{\sqrt{2}}{2} \approx 0.707$$

$$\sqrt{3} \approx 1.732$$

$$\frac{\sqrt{3}}{2} \approx 0.866$$

$$\frac{\sqrt{3}}{3} \approx 0.577$$

1 Find $\sin(\frac{5\pi}{3})$.

Using calculator:

$$\sin(\frac{5\pi}{3}) \approx -0.866$$

$$\Rightarrow \sin(\frac{5\pi}{3}) = -\frac{\sqrt{3}}{2}$$

By hand: Reference angle: $\frac{\pi}{3}$

$$\sin(\frac{5\pi}{3}) = \pm \sin(\frac{\pi}{3}) = \pm \frac{\sqrt{3}}{2}$$

Quadrant of $\frac{5\pi}{3} = 300^\circ$: IV

2 Find $\cos(\frac{5\pi}{4})$.

Calculator:

$$\cos(\frac{5\pi}{4}) \approx -0.707$$

$$\Rightarrow \cos(\frac{5\pi}{4}) = -\frac{\sqrt{2}}{2}$$

Reference angle: $\frac{\pi}{4}$

$$\cos(\frac{5\pi}{4}) = \pm \cos(\frac{\pi}{4}) = \pm \frac{\sqrt{2}}{2}$$

Quadrant of $\frac{5\pi}{4} = 225^\circ$: III

4 Find $\sin(135^\circ)$.

3 Find $\sin(-\frac{11\pi}{6})$.

Calculator:

$$\sin(-\frac{11\pi}{6}) = 0.5$$

$$\Rightarrow \sin(-\frac{11\pi}{6}) = \frac{1}{2}$$

Reference angle: $\frac{\pi}{6}$


$$\sin(-\frac{11\pi}{6}) = \pm \sin(\frac{\pi}{6}) = \pm \frac{1}{2}$$

Quadrant of $-\frac{11\pi}{6} = -330^\circ$: I

5 Find $\cos(210^\circ)$.

Trigonometric functions via reference angles - exercises

Trigonometric functions via reference angle



$$\cos \theta = x$$

$$\sin \theta = y$$

$$\tan \theta = \frac{y}{x}$$

$$\frac{1}{2} = 0.5$$

$$1 = 1$$

$$\frac{\sqrt{2}}{2} \approx 0.707$$

$$\sqrt{3} \approx 1.732$$

$$\frac{\sqrt{3}}{2} \approx 0.866$$

$$\frac{\sqrt{3}}{3} \approx 0.577$$

1 Find $\sin(\frac{5\pi}{3})$.

Using calculator:

$$\sin(\frac{5\pi}{3}) \approx -0.866$$

$$\Rightarrow \sin(\frac{5\pi}{3}) = -\frac{\sqrt{3}}{2}$$

By hand: Reference angle: $\frac{\pi}{3}$

$$\sin(\frac{5\pi}{3}) = \pm \sin(\frac{\pi}{3}) = \pm \frac{\sqrt{3}}{2}$$

$$\text{Quadrant of } \frac{5\pi}{3} = 300^\circ: \text{IV}$$

2 Find $\cos(\frac{5\pi}{4})$.

Calculator:

$$\cos(\frac{5\pi}{4}) \approx -0.707$$

$$\Rightarrow \cos(\frac{5\pi}{4}) = -\frac{\sqrt{2}}{2}$$

Reference angle: $\frac{\pi}{4}$

$$\cos(\frac{5\pi}{4}) = \pm \cos(\frac{\pi}{4}) = \pm \frac{\sqrt{2}}{2}$$

Quadrant of $\frac{5\pi}{4} = 225^\circ$: III

3 Find $\sin(-\frac{11\pi}{6})$.

Calculator:

$$\sin(-\frac{11\pi}{6}) = 0.5$$

$$\Rightarrow \sin(-\frac{11\pi}{6}) = \frac{1}{2}$$

Reference angle: $\frac{\pi}{6}$

$$\sin(-\frac{11\pi}{6}) = \pm \sin(\frac{\pi}{6}) = \pm \frac{1}{2}$$

Quadrant of $-\frac{11\pi}{6} = -330^\circ$: I

4 Find $\sin(135^\circ)$.

Calculator:

$$\sin(135^\circ) \approx 0.707$$

$$\Rightarrow \sin(135^\circ) = \frac{\sqrt{2}}{2}$$

$$135^\circ = 135 \cdot \frac{\pi}{180} = \frac{3\pi}{4}$$

Reference angle: $\frac{\pi}{4}$


$$\sin(\frac{3\pi}{4}) = \pm \sin(\frac{\pi}{4}) = \pm \frac{\sqrt{2}}{2}$$

Quadrant of $\frac{3\pi}{4} = 135^\circ$: II

5 Find $\cos(210^\circ)$.

Trigonometric functions via reference angles - exercises

Trigonometric functions via reference angle



$$\frac{1}{2} = 0.5 \qquad 1 = 1$$

$$\frac{\sqrt{2}}{2} \approx 0.707 \qquad \sqrt{3} \approx 1.732$$

$$\frac{\sqrt{3}}{2} \approx 0.866 \qquad \frac{\sqrt{3}}{3} \approx 0.577$$

1 Find $\sin\left(\frac{5\pi}{3}\right)$.

Using calculator:

$$\sin\left(\frac{5\pi}{3}\right) \approx -0.866$$

$$\Rightarrow \sin\left(\frac{5\pi}{3}\right) = -\frac{\sqrt{3}}{2}$$

By hand: Reference angle: $\frac{\pi}{3}$

$$\sin\left(\frac{5\pi}{3}\right) = \pm \sin\left(\frac{\pi}{3}\right) = \pm \frac{\sqrt{3}}{2}$$

$$\text{Quadrant of } \frac{5\pi}{3} = 300^\circ: \text{ IV}$$

2 Find $\cos\left(\frac{5\pi}{4}\right)$.

Calculator:

$$\cos\left(\frac{5\pi}{4}\right) \approx -0.707$$

$$\Rightarrow \cos\left(\frac{5\pi}{4}\right) = -\frac{\sqrt{2}}{2}$$

Reference angle: $\frac{\pi}{4}$

$$\cos\left(\frac{5\pi}{4}\right) = \pm \cos\left(\frac{\pi}{4}\right) = \pm \frac{\sqrt{2}}{2}$$

Quadrant of $\frac{5\pi}{4} = 225^\circ: \text{ III}$

3 Find $\sin\left(-\frac{11\pi}{6}\right)$.

Calculator:

$$\sin\left(-\frac{11\pi}{6}\right) = 0.5$$

$$\Rightarrow \sin\left(-\frac{11\pi}{6}\right) = \frac{1}{2}$$

Reference angle: $\frac{\pi}{6}$

$$\sin\left(-\frac{11\pi}{6}\right) = \pm \sin\left(\frac{\pi}{6}\right) = \pm \frac{1}{2}$$

Quadrant of $-\frac{11\pi}{6} = -330^\circ: \text{ I}$

4 Find $\sin(135^\circ)$.

Calculator:

$$\sin(135^\circ) \approx 0.707$$

$$\Rightarrow \sin(135^\circ) = \frac{\sqrt{2}}{2}$$

$$135^\circ = 135 \cdot \frac{\pi}{180} = \frac{3\pi}{4}$$

Reference angle: $\frac{\pi}{4}$

$$\sin\left(\frac{3\pi}{4}\right) = \pm \sin\left(\frac{\pi}{4}\right) = \pm \frac{\sqrt{2}}{2}$$

Quadrant of $\frac{3\pi}{4} = 135^\circ: \text{ II}$

5 Find $\cos(210^\circ)$.

Calculator:

$$\cos(210^\circ) \approx -0.866$$

$$\Rightarrow \cos(210^\circ) = -\frac{\sqrt{3}}{2}$$

$$210^\circ = 210 \cdot \frac{\pi}{180} = \frac{7\pi}{6}$$


Reference angle: $\frac{\pi}{6}$

$$\cos\left(\frac{7\pi}{6}\right) = \pm \cos\left(\frac{\pi}{6}\right) = \pm \frac{\sqrt{3}}{2}$$

Quadrant of $\frac{7\pi}{6} = 210^\circ: \text{ III}$

Trigonometric functions via reference angles - exercises

Trigonometric functions via reference angle



$$\cos \theta = x$$

$$\sin \theta = y$$

$$\tan \theta = \frac{y}{x}$$

$$\frac{1}{2} = 0.5$$

$$1 = 1$$

$$\frac{\sqrt{2}}{2} \approx 0.707$$

$$\sqrt{3} \approx 1.732$$

$$\frac{\sqrt{3}}{2} \approx 0.866$$

$$\frac{\sqrt{3}}{3} \approx 0.577$$

6 Find $\tan(\frac{2\pi}{3})$.

7 Find $\tan(\frac{5\pi}{4})$.


9 Find:
 $\sin(\pi)$, $\cos(\pi)$, $\tan(\pi)$

8 Find $\tan(\frac{5\pi}{6})$.

10 Find:
 $\sin(\frac{\pi}{2})$, $\cos(\frac{\pi}{2})$, $\tan(\frac{\pi}{2})$

Trigonometric functions via reference angles - exercises

Trigonometric functions via reference angle



$$\begin{array}{ll} \frac{1}{2} = 0.5 & 1 = 1 \\ \frac{\sqrt{2}}{2} \approx 0.707 & \sqrt{3} \approx 1.732 \\ \frac{\sqrt{3}}{2} \approx 0.866 & \frac{\sqrt{3}}{3} \approx 0.577 \end{array}$$

6 Find $\tan(\frac{2\pi}{3})$.

Using calculator:

$$\tan(\frac{2\pi}{3}) \approx -1.732$$

7 Find $\tan(\frac{5\pi}{4})$.


9 Find:
 $\sin(\pi)$, $\cos(\pi)$, $\tan(\pi)$

8 Find $\tan(\frac{5\pi}{6})$.

10 Find:
 $\sin(\frac{\pi}{2})$, $\cos(\frac{\pi}{2})$, $\tan(\frac{\pi}{2})$

Trigonometric functions via reference angles - exercises

Trigonometric functions via reference angle



$$\cos \theta = x$$

$$\sin \theta = y$$

$$\tan \theta = \frac{y}{x}$$

$$\frac{1}{2} = 0.5$$

$$1 = 1$$

$$\frac{\sqrt{2}}{2} \approx 0.707$$

$$\sqrt{3} \approx 1.732$$

$$\frac{\sqrt{3}}{2} \approx 0.866$$

$$\frac{\sqrt{3}}{3} \approx 0.577$$

6 Find $\tan(\frac{2\pi}{3})$.

Using calculator:

$$\tan(\frac{2\pi}{3}) \approx -1.732$$

$$\Rightarrow \tan(\frac{2\pi}{3}) = -\sqrt{3}$$

7 Find $\tan(\frac{5\pi}{4})$.


9 Find:
 $\sin(\pi)$, $\cos(\pi)$, $\tan(\pi)$

8 Find $\tan(\frac{5\pi}{6})$.

10 Find:
 $\sin(\frac{\pi}{2})$, $\cos(\frac{\pi}{2})$, $\tan(\frac{\pi}{2})$

Trigonometric functions via reference angles - exercises

Trigonometric functions via reference angle



$$\begin{aligned} \frac{1}{2} &= 0.5 & 1 &= 1 \\ \frac{\sqrt{2}}{2} &\approx 0.707 & \sqrt{3} &\approx 1.732 \\ \frac{\sqrt{3}}{2} &\approx 0.866 & \frac{\sqrt{3}}{3} &\approx 0.577 \end{aligned}$$

6 Find $\tan(\frac{2\pi}{3})$.

Using calculator:

$$\tan(\frac{2\pi}{3}) \approx -1.732$$

$$\Rightarrow \tan(\frac{2\pi}{3}) = -\sqrt{3}$$

Reference angle: $\frac{\pi}{3}$

$$\tan(\frac{2\pi}{3}) = \pm \tan(\frac{\pi}{3}) = \pm \frac{\sqrt{3}}{\frac{1}{2}} = \pm \sqrt{3}$$

Quadrant of $\frac{2\pi}{3}$: II

7 Find $\tan(\frac{5\pi}{4})$.


9 Find:
 $\sin(\pi), \cos(\pi), \tan(\pi)$

8 Find $\tan(\frac{5\pi}{6})$.

10 Find:
 $\sin(\frac{\pi}{2}), \cos(\frac{\pi}{2}), \tan(\frac{\pi}{2})$

Trigonometric functions via reference angles - exercises

Trigonometric functions via reference angle



$$\frac{1}{2} = 0.5 \quad 1 = 1$$
$$\frac{\sqrt{2}}{2} \approx 0.707 \quad \sqrt{3} \approx 1.732$$
$$\frac{\sqrt{3}}{2} \approx 0.866 \quad \frac{\sqrt{3}}{3} \approx 0.577$$

6 Find $\tan(\frac{2\pi}{3})$.

Using calculator:

$$\tan(\frac{2\pi}{3}) \approx -1.732$$

$$\Rightarrow \tan(\frac{2\pi}{3}) = -\sqrt{3}$$

Reference angle: $\frac{\pi}{3}$

$$\tan(\frac{2\pi}{3}) = \pm \tan(\frac{\pi}{3}) = \pm \frac{\sqrt{3}}{1} = \pm \sqrt{3}$$

Quadrant of $\frac{2\pi}{3}$: II

7 Find $\tan(\frac{5\pi}{4})$.

Calculator:

$$\tan(\frac{5\pi}{4}) = 1$$

$$\Rightarrow \tan(\frac{5\pi}{4}) = +1$$

Reference angle: $\frac{\pi}{4}$

$$\tan(\frac{5\pi}{4}) = \pm \tan(\frac{\pi}{4})$$
$$= \pm \frac{\frac{\sqrt{2}}{2}}{\frac{\sqrt{2}}{2}} = \pm 1$$

Quadrant of $\frac{5\pi}{4}$: III


8 Find $\tan(\frac{5\pi}{6})$.

9 Find:
 $\sin(\pi)$, $\cos(\pi)$, $\tan(\pi)$

10 Find:
 $\sin(\frac{\pi}{2})$, $\cos(\frac{\pi}{2})$, $\tan(\frac{\pi}{2})$

Trigonometric functions via reference angles - exercises

Trigonometric functions via reference angle



$$\frac{1}{2} = 0.5 \qquad 1 = 1$$

$$\frac{\sqrt{2}}{2} \approx 0.707 \qquad \sqrt{3} \approx 1.732$$

$$\frac{\sqrt{3}}{2} \approx 0.866 \qquad \frac{\sqrt{3}}{3} \approx 0.577$$

6 Find $\tan\left(\frac{2\pi}{3}\right)$.

Using calculator:

$$\tan\left(\frac{2\pi}{3}\right) \approx -1.732$$

$$\Rightarrow \tan\left(\frac{2\pi}{3}\right) = -\sqrt{3}$$

Reference angle: $\frac{\pi}{3}$

$$\tan\left(\frac{2\pi}{3}\right) = \pm \tan\left(\frac{\pi}{3}\right) = \pm \frac{\sqrt{3}}{\frac{1}{2}} = \pm\sqrt{3}$$

Quadrant of $\frac{2\pi}{3}$: II

7 Find $\tan\left(\frac{5\pi}{4}\right)$.

Calculator:

$$\tan\left(\frac{5\pi}{4}\right) = 1$$

$$\Rightarrow \tan\left(\frac{5\pi}{4}\right) = +1$$

Reference angle: $\frac{\pi}{4}$

$$\tan\left(\frac{5\pi}{4}\right) = \pm \tan\left(\frac{\pi}{4}\right)$$

$$= \pm \frac{\frac{\sqrt{2}}{2}}{\frac{\sqrt{2}}{2}} = \pm 1$$

Quadrant of $\frac{5\pi}{4}$: III

8 Find $\tan\left(\frac{5\pi}{6}\right)$.

Calculator:

$$\tan\left(\frac{5\pi}{6}\right) \approx -0.577$$

$$\Rightarrow \tan\left(\frac{5\pi}{6}\right) = -\frac{\sqrt{3}}{3}$$

Reference angle: $\frac{\pi}{6}$

$$\tan\left(\frac{5\pi}{6}\right) = \pm \tan\left(\frac{\pi}{6}\right) = \pm \frac{\frac{1}{2}}{\frac{\sqrt{3}}{2}}$$

$$= \pm \frac{1}{2} \cdot \frac{2}{\sqrt{3}} = \pm \frac{1}{\sqrt{3}} = \pm \frac{\sqrt{3}}{3}$$

Quadrant of $\frac{5\pi}{6}$: II

9 Find:


$$\sin(\pi), \cos(\pi), \tan(\pi)$$

10 Find:

$$\sin\left(\frac{\pi}{2}\right), \cos\left(\frac{\pi}{2}\right), \tan\left(\frac{\pi}{2}\right)$$

Trigonometric functions via reference angles - exercises

Trigonometric functions via reference angle



$$\frac{1}{2} = 0.5 \qquad 1 = 1$$

$$\frac{\sqrt{2}}{2} \approx 0.707 \qquad \sqrt{3} \approx 1.732$$

$$\frac{\sqrt{3}}{2} \approx 0.866 \qquad \frac{\sqrt{3}}{3} \approx 0.577$$

6 Find $\tan(\frac{2\pi}{3})$.

Using calculator:

$$\tan(\frac{2\pi}{3}) \approx -1.732$$

$$\Rightarrow \tan(\frac{2\pi}{3}) = -\sqrt{3}$$

Reference angle: $\frac{\pi}{3}$

$$\tan(\frac{2\pi}{3}) = \pm \tan(\frac{\pi}{3}) = \pm \frac{\sqrt{3}}{\frac{1}{2}} = \pm \sqrt{3}$$

Quadrant of $\frac{2\pi}{3}$: II

7 Find $\tan(\frac{5\pi}{4})$.

Calculator:

$$\tan(\frac{5\pi}{4}) = 1$$

$$\Rightarrow \tan(\frac{5\pi}{4}) = +1$$

Reference angle: $\frac{\pi}{4}$

$$\tan(\frac{5\pi}{4}) = \pm \tan(\frac{\pi}{4})$$

$$= \pm \frac{\frac{\sqrt{2}}{2}}{\frac{\sqrt{2}}{2}} = \pm 1$$

Quadrant of $\frac{5\pi}{4}$: III

8 Find $\tan(\frac{5\pi}{6})$.

Calculator:

$$\tan(\frac{5\pi}{6}) \approx -0.577$$

$$\Rightarrow \tan(\frac{5\pi}{6}) = -\frac{\sqrt{3}}{3}$$

Reference angle: $\frac{\pi}{6}$


$$\tan(\frac{5\pi}{6}) = \pm \tan(\frac{\pi}{6}) = \pm \frac{\frac{1}{2}}{\frac{\sqrt{3}}{2}}$$

$$= \pm \frac{1}{2} \cdot \frac{2}{\sqrt{3}} = \pm \frac{1}{\sqrt{3}} = \pm \frac{\sqrt{3}}{3}$$

Quadrant of $\frac{5\pi}{6}$: II

9 Find:

$$\sin(\pi), \cos(\pi), \tan(\pi)$$



$$\Rightarrow \sin(\pi) = 0$$

$$\Rightarrow \cos(\pi) = -1$$

$$\Rightarrow \tan(\pi) = \frac{0}{-1} = 0$$


(Check with calculator)

10 Find:

$$\sin(\frac{\pi}{2}), \cos(\frac{\pi}{2}), \tan(\frac{\pi}{2})$$

Trigonometric functions via reference angles - exercises

Trigonometric functions via reference angle



$$\frac{1}{2} = 0.5 \qquad 1 = 1$$

$$\frac{\sqrt{2}}{2} \approx 0.707 \qquad \sqrt{3} \approx 1.732$$

$$\frac{\sqrt{3}}{2} \approx 0.866 \qquad \frac{\sqrt{3}}{3} \approx 0.577$$

6 Find $\tan(\frac{2\pi}{3})$.

Using calculator:

$$\tan(\frac{2\pi}{3}) \approx -1.732$$

$$\Rightarrow \tan(\frac{2\pi}{3}) = -\sqrt{3}$$

Reference angle: $\frac{\pi}{3}$

$$\tan(\frac{2\pi}{3}) = \pm \tan(\frac{\pi}{3}) = \pm \frac{\sqrt{3}}{\frac{1}{2}} = \pm \sqrt{3}$$

Quadrant of $\frac{2\pi}{3}$: II

7 Find $\tan(\frac{5\pi}{4})$.

Calculator:

$$\tan(\frac{5\pi}{4}) = 1$$

$$\Rightarrow \tan(\frac{5\pi}{4}) = +1$$

Reference angle: $\frac{\pi}{4}$

$$\tan(\frac{5\pi}{4}) = \pm \tan(\frac{\pi}{4})$$

$$= \pm \frac{\frac{\sqrt{2}}{2}}{\frac{\sqrt{2}}{2}} = \pm 1$$

Quadrant of $\frac{5\pi}{4}$: III

8 Find $\tan(\frac{5\pi}{6})$.

Calculator:

$$\tan(\frac{5\pi}{6}) \approx -0.577$$

$$\Rightarrow \tan(\frac{5\pi}{6}) = -\frac{\sqrt{3}}{3}$$

Reference angle: $\frac{\pi}{6}$


$$\tan(\frac{5\pi}{6}) = \pm \tan(\frac{\pi}{6}) = \pm \frac{\frac{1}{\sqrt{3}}}{\frac{\sqrt{3}}{2}}$$

$$= \pm \frac{1}{2} \cdot \frac{2}{\sqrt{3}} = \pm \frac{1}{\sqrt{3}} = \pm \frac{\sqrt{3}}{3}$$

Quadrant of $\frac{5\pi}{6}$: II

9 Find:

$$\sin(\pi), \cos(\pi), \tan(\pi)$$



$$\Rightarrow \sin(\pi) = 0$$


$$\Rightarrow \cos(\pi) = -1$$

$$\Rightarrow \tan(\pi) = \frac{0}{-1} = 0$$

(Check with calculator)

10 Find:

$$\sin(\frac{\pi}{2}), \cos(\frac{\pi}{2}), \tan(\frac{\pi}{2})$$



$$\Rightarrow \sin(\frac{\pi}{2}) = 1$$

$$\Rightarrow \cos(\frac{\pi}{2}) = 0$$

$$\Rightarrow \tan(\frac{\pi}{2}) = \frac{1}{0} = \text{"1/0" undefined}$$

(Check with calculator)

Sum and difference formulas

Sum and difference formulas

$$\sin(\alpha + \beta) = \sin(\alpha) \cos(\beta) + \cos(\alpha) \sin(\beta)$$

$$\sin(\alpha - \beta) = \sin(\alpha) \cos(\beta) - \cos(\alpha) \sin(\beta)$$

$$\cos(\alpha + \beta) = \cos(\alpha) \cos(\beta) - \sin(\alpha) \sin(\beta)$$

$$\cos(\alpha - \beta) = \cos(\alpha) \cos(\beta) + \sin(\alpha) \sin(\beta)$$

$$\tan(\alpha + \beta) = \frac{\tan(\alpha) + \tan(\beta)}{1 - \tan(\alpha) \tan(\beta)}$$

$$\tan(\alpha - \beta) = \frac{\tan(\alpha) - \tan(\beta)}{1 + \tan(\alpha) \tan(\beta)}$$

1 Find $\cos(105^\circ)$.

2 Find $\sin(15^\circ)$.

3 Find $\tan\left(\frac{19\pi}{12}\right)$.

Sum and difference formulas

Sum and difference formulas

$$\sin(\alpha + \beta) = \sin(\alpha) \cos(\beta) + \cos(\alpha) \sin(\beta)$$

$$\sin(\alpha - \beta) = \sin(\alpha) \cos(\beta) - \cos(\alpha) \sin(\beta)$$

$$\cos(\alpha + \beta) = \cos(\alpha) \cos(\beta) - \sin(\alpha) \sin(\beta)$$

$$\cos(\alpha - \beta) = \cos(\alpha) \cos(\beta) + \sin(\alpha) \sin(\beta)$$

$$\tan(\alpha + \beta) = \frac{\tan(\alpha) + \tan(\beta)}{1 - \tan(\alpha) \tan(\beta)}$$

$$\tan(\alpha - \beta) = \frac{\tan(\alpha) - \tan(\beta)}{1 + \tan(\alpha) \tan(\beta)}$$

- 1 Find $\cos(105^\circ)$.

Note: $105^\circ = 60^\circ + 45^\circ$

- 2 Find $\sin(15^\circ)$.

- 3 Find $\tan\left(\frac{19\pi}{12}\right)$.

Sum and difference formulas

Sum and difference formulas

$$\sin(\alpha + \beta) = \sin(\alpha) \cos(\beta) + \cos(\alpha) \sin(\beta)$$

$$\sin(\alpha - \beta) = \sin(\alpha) \cos(\beta) - \cos(\alpha) \sin(\beta)$$

$$\cos(\alpha + \beta) = \cos(\alpha) \cos(\beta) - \sin(\alpha) \sin(\beta)$$

$$\cos(\alpha - \beta) = \cos(\alpha) \cos(\beta) + \sin(\alpha) \sin(\beta)$$

$$\tan(\alpha + \beta) = \frac{\tan(\alpha) + \tan(\beta)}{1 - \tan(\alpha) \tan(\beta)}$$

$$\tan(\alpha - \beta) = \frac{\tan(\alpha) - \tan(\beta)}{1 + \tan(\alpha) \tan(\beta)}$$

- 1 Find $\cos(105^\circ)$.

Note: $105^\circ = 60^\circ + 45^\circ$

$$\cos(105^\circ) = \cos(60^\circ + 45^\circ)$$

$$= \cos(60^\circ) \cos(45^\circ) - \sin(60^\circ) \sin(45^\circ)$$

$$= \frac{1}{2} \cdot \frac{\sqrt{2}}{2} - \frac{\sqrt{3}}{2} \cdot \frac{\sqrt{2}}{2}$$

$$= \frac{\sqrt{2}}{4} - \frac{\sqrt{3} \cdot \sqrt{2}}{4} = \frac{\sqrt{2} - \sqrt{6}}{4}$$

- 2 Find $\sin(15^\circ)$.

- 3 Find $\tan\left(\frac{19\pi}{12}\right)$.

Sum and difference formulas

Sum and difference formulas

$$\sin(\alpha + \beta) = \sin(\alpha) \cos(\beta) + \cos(\alpha) \sin(\beta)$$

$$\sin(\alpha - \beta) = \sin(\alpha) \cos(\beta) - \cos(\alpha) \sin(\beta)$$

$$\cos(\alpha + \beta) = \cos(\alpha) \cos(\beta) - \sin(\alpha) \sin(\beta)$$

$$\cos(\alpha - \beta) = \cos(\alpha) \cos(\beta) + \sin(\alpha) \sin(\beta)$$

$$\tan(\alpha + \beta) = \frac{\tan(\alpha) + \tan(\beta)}{1 - \tan(\alpha) \tan(\beta)}$$

$$\tan(\alpha - \beta) = \frac{\tan(\alpha) - \tan(\beta)}{1 + \tan(\alpha) \tan(\beta)}$$

- 1 Find $\cos(105^\circ)$.

Note: $105^\circ = 60^\circ + 45^\circ$

$$\cos(105^\circ) = \cos(60^\circ + 45^\circ)$$

$$= \cos(60^\circ) \cos(45^\circ) - \sin(60^\circ) \sin(45^\circ)$$

$$= \frac{1}{2} \cdot \frac{\sqrt{2}}{2} - \frac{\sqrt{3}}{2} \cdot \frac{\sqrt{2}}{2}$$

$$= \frac{\sqrt{2}}{4} - \frac{\sqrt{3} \cdot \sqrt{2}}{4} = \frac{\sqrt{2} - \sqrt{6}}{4}$$

- 2 Find $\sin(15^\circ)$.

Note: $15^\circ = 60^\circ - 45^\circ$

- 3 Find $\tan\left(\frac{19\pi}{12}\right)$.

Sum and difference formulas

Sum and difference formulas

$$\sin(\alpha + \beta) = \sin(\alpha) \cos(\beta) + \cos(\alpha) \sin(\beta)$$

$$\sin(\alpha - \beta) = \sin(\alpha) \cos(\beta) - \cos(\alpha) \sin(\beta)$$

$$\cos(\alpha + \beta) = \cos(\alpha) \cos(\beta) - \sin(\alpha) \sin(\beta)$$

$$\cos(\alpha - \beta) = \cos(\alpha) \cos(\beta) + \sin(\alpha) \sin(\beta)$$

$$\tan(\alpha + \beta) = \frac{\tan(\alpha) + \tan(\beta)}{1 - \tan(\alpha) \tan(\beta)}$$

$$\tan(\alpha - \beta) = \frac{\tan(\alpha) - \tan(\beta)}{1 + \tan(\alpha) \tan(\beta)}$$

- 1 Find $\cos(105^\circ)$.

Note: $105^\circ = 60^\circ + 45^\circ$

$$\cos(105^\circ) = \cos(60^\circ + 45^\circ)$$

$$= \cos(60^\circ) \cos(45^\circ) - \sin(60^\circ) \sin(45^\circ)$$

$$= \frac{1}{2} \cdot \frac{\sqrt{2}}{2} - \frac{\sqrt{3}}{2} \cdot \frac{\sqrt{2}}{2}$$

$$= \frac{\sqrt{2}}{4} - \frac{\sqrt{3} \cdot \sqrt{2}}{4} = \frac{\sqrt{2} - \sqrt{6}}{4}$$

- 2 Find $\sin(15^\circ)$.

Note: $15^\circ = 60^\circ - 45^\circ$

$$\sin(15^\circ) = \sin(60^\circ - 45^\circ)$$

$$= \sin(60^\circ) \cos(45^\circ) - \cos(60^\circ) \sin(45^\circ)$$

$$= \frac{\sqrt{3}}{2} \cdot \frac{\sqrt{2}}{2} - \frac{1}{2} \cdot \frac{\sqrt{2}}{2}$$

$$= \frac{\sqrt{3} \cdot \sqrt{2}}{4} - \frac{\sqrt{2}}{4}$$

$$= \frac{\sqrt{6} - \sqrt{2}}{4}$$

- 3 Find $\tan\left(\frac{19\pi}{12}\right)$.

Sum and difference formulas

Sum and difference formulas

$$\sin(\alpha + \beta) = \sin(\alpha) \cos(\beta) + \cos(\alpha) \sin(\beta)$$

$$\sin(\alpha - \beta) = \sin(\alpha) \cos(\beta) - \cos(\alpha) \sin(\beta)$$

$$\cos(\alpha + \beta) = \cos(\alpha) \cos(\beta) - \sin(\alpha) \sin(\beta)$$

$$\cos(\alpha - \beta) = \cos(\alpha) \cos(\beta) + \sin(\alpha) \sin(\beta)$$

$$\tan(\alpha + \beta) = \frac{\tan(\alpha) + \tan(\beta)}{1 - \tan(\alpha) \tan(\beta)}$$

$$\tan(\alpha - \beta) = \frac{\tan(\alpha) - \tan(\beta)}{1 + \tan(\alpha) \tan(\beta)}$$

- 1 Find $\cos(105^\circ)$.

Note: $105^\circ = 60^\circ + 45^\circ$

$$\cos(105^\circ) = \cos(60^\circ + 45^\circ)$$

$$= \cos(60^\circ) \cos(45^\circ) - \sin(60^\circ) \sin(45^\circ)$$

$$= \frac{1}{2} \cdot \frac{\sqrt{2}}{2} - \frac{\sqrt{3}}{2} \cdot \frac{\sqrt{2}}{2}$$

$$= \frac{\sqrt{2}}{4} - \frac{\sqrt{3} \cdot \sqrt{2}}{4} = \frac{\sqrt{2} - \sqrt{6}}{4}$$

- 2 Find $\sin(15^\circ)$.

Note: $15^\circ = 60^\circ - 45^\circ$

$$\sin(15^\circ) = \sin(60^\circ - 45^\circ)$$

$$= \sin(60^\circ) \cos(45^\circ) - \cos(60^\circ) \sin(45^\circ)$$

$$= \frac{\sqrt{3}}{2} \cdot \frac{\sqrt{2}}{2} - \frac{1}{2} \cdot \frac{\sqrt{2}}{2}$$

$$= \frac{\sqrt{3} \cdot \sqrt{2}}{4} - \frac{\sqrt{2}}{4}$$

$$= \frac{\sqrt{6} - \sqrt{2}}{4}$$

- 3 Find $\tan\left(\frac{19\pi}{12}\right)$.

Note: $\frac{19\pi}{12} = 285^\circ = 60^\circ + 225^\circ$

Sum and difference formulas

Sum and difference formulas

$$\sin(\alpha + \beta) = \sin(\alpha) \cos(\beta) + \cos(\alpha) \sin(\beta)$$

$$\sin(\alpha - \beta) = \sin(\alpha) \cos(\beta) - \cos(\alpha) \sin(\beta)$$

$$\cos(\alpha + \beta) = \cos(\alpha) \cos(\beta) - \sin(\alpha) \sin(\beta)$$

$$\cos(\alpha - \beta) = \cos(\alpha) \cos(\beta) + \sin(\alpha) \sin(\beta)$$

$$\tan(\alpha + \beta) = \frac{\tan(\alpha) + \tan(\beta)}{1 - \tan(\alpha) \tan(\beta)}$$

$$\tan(\alpha - \beta) = \frac{\tan(\alpha) - \tan(\beta)}{1 + \tan(\alpha) \tan(\beta)}$$

- 1 Find $\cos(105^\circ)$.

Note: $105^\circ = 60^\circ + 45^\circ$

$$\cos(105^\circ) = \cos(60^\circ + 45^\circ)$$

$$= \cos(60^\circ) \cos(45^\circ) - \sin(60^\circ) \sin(45^\circ)$$

$$= \frac{1}{2} \cdot \frac{\sqrt{2}}{2} - \frac{\sqrt{3}}{2} \cdot \frac{\sqrt{2}}{2}$$

$$= \frac{\sqrt{2}}{4} - \frac{\sqrt{3} \cdot \sqrt{2}}{4} = \frac{\sqrt{2} - \sqrt{6}}{4}$$

- 2 Find $\sin(15^\circ)$.

Note: $15^\circ = 60^\circ - 45^\circ$

$$\sin(15^\circ) = \sin(60^\circ - 45^\circ)$$

$$= \sin(60^\circ) \cos(45^\circ) - \cos(60^\circ) \sin(45^\circ)$$

$$= \frac{\sqrt{3}}{2} \cdot \frac{\sqrt{2}}{2} - \frac{1}{2} \cdot \frac{\sqrt{2}}{2}$$

$$= \frac{\sqrt{3} \cdot \sqrt{2}}{4} - \frac{\sqrt{2}}{4}$$

$$= \frac{\sqrt{6} - \sqrt{2}}{4}$$

- 3 Find $\tan\left(\frac{19\pi}{12}\right)$.

Note: $\frac{19\pi}{12} = 285^\circ = 60^\circ + 225^\circ$

$$\tan\left(\frac{19\pi}{12}\right) = \tan(285^\circ)$$

$$= \tan(60^\circ + 225^\circ)$$

Sum and difference formulas

Sum and difference formulas

$$\sin(\alpha + \beta) = \sin(\alpha) \cos(\beta) + \cos(\alpha) \sin(\beta)$$

$$\sin(\alpha - \beta) = \sin(\alpha) \cos(\beta) - \cos(\alpha) \sin(\beta)$$

$$\cos(\alpha + \beta) = \cos(\alpha) \cos(\beta) - \sin(\alpha) \sin(\beta)$$

$$\cos(\alpha - \beta) = \cos(\alpha) \cos(\beta) + \sin(\alpha) \sin(\beta)$$

$$\tan(\alpha + \beta) = \frac{\tan(\alpha) + \tan(\beta)}{1 - \tan(\alpha) \tan(\beta)}$$

$$\tan(\alpha - \beta) = \frac{\tan(\alpha) - \tan(\beta)}{1 + \tan(\alpha) \tan(\beta)}$$

- 1 Find $\cos(105^\circ)$.

Note: $105^\circ = 60^\circ + 45^\circ$

$$\cos(105^\circ) = \cos(60^\circ + 45^\circ)$$

$$= \cos(60^\circ) \cos(45^\circ) - \sin(60^\circ) \sin(45^\circ)$$

$$= \frac{1}{2} \cdot \frac{\sqrt{2}}{2} - \frac{\sqrt{3}}{2} \cdot \frac{\sqrt{2}}{2}$$

$$= \frac{\sqrt{2}}{4} - \frac{\sqrt{3} \cdot \sqrt{2}}{4} = \frac{\sqrt{2} - \sqrt{6}}{4}$$

- 2 Find $\sin(15^\circ)$.

Note: $15^\circ = 60^\circ - 45^\circ$

$$\sin(15^\circ) = \sin(60^\circ - 45^\circ)$$

$$= \sin(60^\circ) \cos(45^\circ) - \cos(60^\circ) \sin(45^\circ)$$

$$= \frac{\sqrt{3}}{2} \cdot \frac{\sqrt{2}}{2} - \frac{1}{2} \cdot \frac{\sqrt{2}}{2}$$

$$= \frac{\sqrt{3} \cdot \sqrt{2}}{4} - \frac{\sqrt{2}}{4}$$

$$= \frac{\sqrt{6} - \sqrt{2}}{4}$$

- 3 Find $\tan\left(\frac{19\pi}{12}\right)$.

Note: $\frac{19\pi}{12} = 285^\circ = 60^\circ + 225^\circ$

$$\tan\left(\frac{19\pi}{12}\right) = \tan(285^\circ)$$

$$= \tan(60^\circ + 225^\circ)$$

$$= \frac{\tan(60^\circ) + \tan(225^\circ)}{1 - \tan(60^\circ) \cdot \tan(225^\circ)}$$

$$= \frac{\sqrt{3} + 1}{1 - \sqrt{3} \cdot 1} = \frac{1 + \sqrt{3}}{1 - \sqrt{3}}$$

Double and half angle formulas

Half angle formulas

$$\sin\left(\frac{\alpha}{2}\right) = \pm\sqrt{\frac{1 - \cos(\alpha)}{2}}$$

$$\cos\left(\frac{\alpha}{2}\right) = \pm\sqrt{\frac{1 + \cos(\alpha)}{2}}$$

$$\tan\left(\frac{\alpha}{2}\right) = \pm\sqrt{\frac{1 - \cos(\alpha)}{1 + \cos(\alpha)}}$$

Double angle formulas

$$\sin(2\alpha) = 2 \sin(\alpha) \cos(\alpha)$$

$$\begin{aligned}\cos(2\alpha) &= \cos^2(\alpha) - \sin^2(\alpha) \\ &= 1 - 2 \sin^2(\alpha) = 2 \cos^2(\alpha) - 1\end{aligned}$$

$$\tan(2\alpha) = \frac{2 \tan(\alpha)}{1 - \tan^2(\alpha)}$$

Double and half angle formulas

Half angle formulas

$$\sin\left(\frac{\alpha}{2}\right) = \pm\sqrt{\frac{1 - \cos(\alpha)}{2}}$$

$$\cos\left(\frac{\alpha}{2}\right) = \pm\sqrt{\frac{1 + \cos(\alpha)}{2}}$$

$$\tan\left(\frac{\alpha}{2}\right) = \pm\sqrt{\frac{1 - \cos(\alpha)}{1 + \cos(\alpha)}}$$

- 1 Find $\sin(15^\circ)$
using the half angle formulas.

Double angle formulas

$$\sin(2\alpha) = 2 \sin(\alpha) \cos(\alpha)$$

$$\begin{aligned}\cos(2\alpha) &= \cos^2(\alpha) - \sin^2(\alpha) \\ &= 1 - 2 \sin^2(\alpha) = 2 \cos^2(\alpha) - 1\end{aligned}$$

$$\tan(2\alpha) = \frac{2 \tan(\alpha)}{1 - \tan^2(\alpha)}$$

- 2 Find $\cos\left(\frac{\pi}{8}\right)$
using the half angle formulas.

Double and half angle formulas

Half angle formulas

$$\sin\left(\frac{\alpha}{2}\right) = \pm \sqrt{\frac{1 - \cos(\alpha)}{2}}$$

$$\cos\left(\frac{\alpha}{2}\right) = \pm \sqrt{\frac{1 + \cos(\alpha)}{2}}$$

$$\tan\left(\frac{\alpha}{2}\right) = \pm \sqrt{\frac{1 - \cos(\alpha)}{1 + \cos(\alpha)}}$$

- 1 Find $\sin(15^\circ)$ using the half angle formulas.

Note: $15^\circ = \frac{30^\circ}{2}$

Double angle formulas

$$\sin(2\alpha) = 2 \sin(\alpha) \cos(\alpha)$$

$$\begin{aligned} \cos(2\alpha) &= \cos^2(\alpha) - \sin^2(\alpha) \\ &= 1 - 2 \sin^2(\alpha) = 2 \cos^2(\alpha) - 1 \end{aligned}$$

$$\tan(2\alpha) = \frac{2 \tan(\alpha)}{1 - \tan^2(\alpha)}$$

- 2 Find $\cos\left(\frac{\pi}{8}\right)$ using the half angle formulas.

Double and half angle formulas

Half angle formulas

$$\sin\left(\frac{\alpha}{2}\right) = \pm \sqrt{\frac{1 - \cos(\alpha)}{2}}$$

$$\cos\left(\frac{\alpha}{2}\right) = \pm \sqrt{\frac{1 + \cos(\alpha)}{2}}$$

$$\tan\left(\frac{\alpha}{2}\right) = \pm \sqrt{\frac{1 - \cos(\alpha)}{1 + \cos(\alpha)}}$$

Double angle formulas

$$\sin(2\alpha) = 2 \sin(\alpha) \cos(\alpha)$$

$$\begin{aligned} \cos(2\alpha) &= \cos^2(\alpha) - \sin^2(\alpha) \\ &= 1 - 2 \sin^2(\alpha) = 2 \cos^2(\alpha) - 1 \end{aligned}$$

$$\tan(2\alpha) = \frac{2 \tan(\alpha)}{1 - \tan^2(\alpha)}$$

- ① Find $\sin(15^\circ)$ using the half angle formulas.

Note: $15^\circ = \frac{30^\circ}{2}$

$$\implies \sin(15^\circ) = \sin\left(\frac{30^\circ}{2}\right)$$

$$= + \sqrt{\frac{1 - \cos(30^\circ)}{2}} = \sqrt{\frac{1 - \frac{\sqrt{3}}{2}}{2}}$$

$$= \sqrt{\frac{\frac{2 - \sqrt{3}}{2}}{2}} = \sqrt{\frac{2 - \sqrt{3}}{2}} \cdot \frac{1}{2}$$

$$= \sqrt{\frac{2 - \sqrt{3}}{4}} = \frac{\sqrt{2 - \sqrt{3}}}{2}$$

- ② Find $\cos\left(\frac{\pi}{8}\right)$ using the half angle formulas.

Double and half angle formulas

Half angle formulas

$$\sin\left(\frac{\alpha}{2}\right) = \pm \sqrt{\frac{1 - \cos(\alpha)}{2}}$$

$$\cos\left(\frac{\alpha}{2}\right) = \pm \sqrt{\frac{1 + \cos(\alpha)}{2}}$$

$$\tan\left(\frac{\alpha}{2}\right) = \pm \sqrt{\frac{1 - \cos(\alpha)}{1 + \cos(\alpha)}}$$

Double angle formulas

$$\sin(2\alpha) = 2 \sin(\alpha) \cos(\alpha)$$

$$\begin{aligned} \cos(2\alpha) &= \cos^2(\alpha) - \sin^2(\alpha) \\ &= 1 - 2 \sin^2(\alpha) = 2 \cos^2(\alpha) - 1 \end{aligned}$$

$$\tan(2\alpha) = \frac{2 \tan(\alpha)}{1 - \tan^2(\alpha)}$$

- 1 Find $\sin(15^\circ)$ using the half angle formulas.

Note: $15^\circ = \frac{30^\circ}{2}$

$$\implies \sin(15^\circ) = \sin\left(\frac{30^\circ}{2}\right)$$

$$= + \sqrt{\frac{1 - \cos(30^\circ)}{2}} = \sqrt{\frac{1 - \frac{\sqrt{3}}{2}}{2}}$$

$$= \sqrt{\frac{\frac{2 - \sqrt{3}}{2}}{2}} = \sqrt{\frac{2 - \sqrt{3}}{2}} \cdot \frac{1}{2}$$

$$= \sqrt{\frac{2 - \sqrt{3}}{4}} = \frac{\sqrt{2 - \sqrt{3}}}{2}$$

- 2 Find $\cos\left(\frac{\pi}{8}\right)$ using the half angle formulas.

Note: $\frac{\pi}{8} = 22.5^\circ = \frac{45^\circ}{2}$

Double and half angle formulas

Half angle formulas

$$\sin\left(\frac{\alpha}{2}\right) = \pm \sqrt{\frac{1 - \cos(\alpha)}{2}}$$

$$\cos\left(\frac{\alpha}{2}\right) = \pm \sqrt{\frac{1 + \cos(\alpha)}{2}}$$

$$\tan\left(\frac{\alpha}{2}\right) = \pm \sqrt{\frac{1 - \cos(\alpha)}{1 + \cos(\alpha)}}$$

Double angle formulas

$$\sin(2\alpha) = 2 \sin(\alpha) \cos(\alpha)$$

$$\begin{aligned}\cos(2\alpha) &= \cos^2(\alpha) - \sin^2(\alpha) \\ &= 1 - 2 \sin^2(\alpha) = 2 \cos^2(\alpha) - 1\end{aligned}$$

$$\tan(2\alpha) = \frac{2 \tan(\alpha)}{1 - \tan^2(\alpha)}$$

- 1 Find $\sin(15^\circ)$ using the half angle formulas.

Note: $15^\circ = \frac{30^\circ}{2}$

$$\implies \sin(15^\circ) = \sin\left(\frac{30^\circ}{2}\right)$$

$$= + \sqrt{\frac{1 - \cos(30^\circ)}{2}} = \sqrt{\frac{1 - \frac{\sqrt{3}}{2}}{2}}$$

$$= \sqrt{\frac{\frac{2 - \sqrt{3}}{2}}{2}} = \sqrt{\frac{2 - \sqrt{3}}{2}} \cdot \frac{1}{2}$$

$$= \sqrt{\frac{2 - \sqrt{3}}{4}} = \frac{\sqrt{2 - \sqrt{3}}}{2}$$

- 2 Find $\cos\left(\frac{\pi}{8}\right)$ using the half angle formulas.

Note: $\frac{\pi}{8} = 22.5^\circ = \frac{45^\circ}{2}$

$$\implies \cos\left(\frac{\pi}{8}\right) = \cos\left(\frac{45^\circ}{2}\right)$$

$$= + \sqrt{\frac{1 + \cos(45^\circ)}{2}} = \sqrt{\frac{1 + \frac{\sqrt{2}}{2}}{2}}$$

$$= \sqrt{\frac{\frac{2 + \sqrt{2}}{2}}{2}} = \sqrt{\frac{2 + \sqrt{2}}{2}} \cdot \frac{1}{2}$$

$$= \sqrt{\frac{2 + \sqrt{2}}{4}} = \frac{\sqrt{2 + \sqrt{2}}}{2}$$

