

Geometric series

Lesson #25

MAT 1375 Precalculus

New York City College of Technology CUNY



Geometric sequences

Geometric sequence

An **geometric sequence** is a sequence where each term follows from its previous term by multiplying a fixed number r :


$$a_1 \xrightarrow{\cdot r} a_2 \xrightarrow{\cdot r} a_3 \xrightarrow{\cdot r} a_4 \xrightarrow{\cdot r} a_5 \dots$$

r is called the **common ratio**.

Geometric sequences

Geometric sequence

An **geometric sequence** is a sequence where each term follows from its previous term by multiplying a fixed number r :



r is called the **common ratio**.


Example

- Find the first 5 terms of the sequence $a_n = 3 \cdot 10^{n-1}$.

Geometric sequences

Geometric sequence

An **geometric sequence** is a sequence where each term follows from its previous term by multiplying a fixed number r :



r is called the **common ratio**.

Example

- ① Find the first 5 terms of the sequence $a_n = 3 \cdot 10^{n-1}$.

Answer:


3, 30, 300, 3000, 30000, ...

This is a **geometric sequence** with first term $a_1 = 3$ and common ratio $r = 10$.

Geometric sequences

Geometric sequence

An **geometric sequence** is a sequence where each term follows from its previous term by multiplying a fixed number r :



r is called the **common ratio**.

Example

- ① Find the first 5 terms of the sequence $a_n = 3 \cdot 10^{n-1}$.

Answer:

3, 30, 300, 3000, 30000, ...


This is a **geometric sequence** with first term $a_1 = 3$ and common ratio $r = 10$.

- ② Find the rule of the geometric sequence: 5, 10, 20, 40, 80, 160, ...

Geometric sequences

Geometric sequence

An **geometric sequence** is a sequence where each term follows from its previous term by multiplying a fixed number r :



r is called the **common ratio**.

Example

- ① Find the first 5 terms of the sequence $a_n = 3 \cdot 10^{n-1}$.

Answer:

$$3, 30, 300, 3000, 30000, \dots$$

This is a **geometric sequence** with first term $a_1 = 3$ and common ratio $r = 10$.

- ② Find the rule of the geometric sequence: 5, 10, 20, 40, 80, 160, ...


Answer:

$$a_n = 5 \cdot 2^{n-1}$$

Geometric sequences

Geometric sequence

An **geometric sequence** is a sequence where each term follows from its previous term by multiplying a fixed number r :



r is called the **common ratio**. A geometric sequence has the closed form formula:

$$a_n = a_1 \cdot r^{n-1}$$

Example

- ① Find the first 5 terms of the sequence $a_n = 3 \cdot 10^{n-1}$.

Answer:

3, 30, 300, 3000, 30000, ...

This is a **geometric sequence** with first term $a_1 = 3$ and common ratio $r = 10$.

- ② Find the rule of the geometric sequence: 5, 10, 20, 40, 80, 160, ...

Answer:

$$a_n = 5 \cdot 2^{n-1}$$

Geometric sequence

$$a_n = a_1 \cdot r^{n-1}$$

Find the closed form formula for the given geometric sequence.

- 1 7, 14, 28, 56, ...
- 2 4, -20, 100, -500, ...
- 3 8, 0.8, 0.08, 0.008, ...
- 4 48, 24, 12, 6, 3, $\frac{3}{2}$, $\frac{3}{4}$, ...

Geometric sequence

$$a_n = a_1 \cdot r^{n-1}$$

Find the closed form formula for the given geometric sequence.

① 7, 14, 28, 56, ...

$$a_1 = 7, r = 2, \Rightarrow a_n = 7 \cdot 2^{n-1}$$

② 4, -20, 100, -500, ...

③ 8, 0.8, 0.08, 0.008, ...

④ 48, 24, 12, 6, 3, $\frac{3}{2}$, $\frac{3}{4}$, ...

Geometric sequence

$$a_n = a_1 \cdot r^{n-1}$$

Find the closed form formula for the given geometric sequence.

① 7, 14, 28, 56, ...

$$a_1 = 7, r = 2, \Rightarrow a_n = 7 \cdot 2^{n-1}$$

② 4, -20, 100, -500, ...

$$a_1 = 4, r = -5, \Rightarrow a_n = 4 \cdot (-5)^{n-1}$$

③ 8, 0.8, 0.08, 0.008, ...

④ 48, 24, 12, 6, 3, $\frac{3}{2}$, $\frac{3}{4}$, ...

Geometric sequence

$$a_n = a_1 \cdot r^{n-1}$$

Find the closed form formula for the given geometric sequence.

① 7, 14, 28, 56, ...

$$a_1 = 7, r = 2, \Rightarrow a_n = 7 \cdot 2^{n-1}$$

② 4, -20, 100, -500, ...

$$a_1 = 4, r = -5, \Rightarrow a_n = 4 \cdot (-5)^{n-1}$$

③ 8, 0.8, 0.08, 0.008, ...

$$a_1 = 8, r = \frac{1}{10}, \Rightarrow a_n = 8 \cdot \left(\frac{1}{10}\right)^{n-1}$$

④ 48, 24, 12, 6, 3, $\frac{3}{2}$, $\frac{3}{4}$, ...

Geometric sequence

$$a_n = a_1 \cdot r^{n-1}$$

Find the closed form formula for the given geometric sequence.

① 7, 14, 28, 56, ...

$$a_1 = 7, r = 2, \Rightarrow a_n = 7 \cdot 2^{n-1}$$

② 4, -20, 100, -500, ...

$$a_1 = 4, r = -5, \Rightarrow a_n = 4 \cdot (-5)^{n-1}$$

③ 8, 0.8, 0.08, 0.008, ...

$$a_1 = 8, r = \frac{1}{10}, \Rightarrow a_n = 8 \cdot \left(\frac{1}{10}\right)^{n-1}$$

④ 48, 24, 12, 6, 3, $\frac{3}{2}$, $\frac{3}{4}$, ...

$$a_1 = 48, r = \frac{1}{2}, \Rightarrow a_n = 48 \cdot \left(\frac{1}{2}\right)^{n-1}$$

Geometric sequence

$$a_n = a_1 \cdot r^{n-1}$$

Find the closed form formula for the given geometric sequence.

① 7, 14, 28, 56, ...

$$a_1 = 7, r = 2, \Rightarrow a_n = 7 \cdot 2^{n-1}$$

② 4, -20, 100, -500, ...

$$a_1 = 4, r = -5, \Rightarrow a_n = 4 \cdot (-5)^{n-1}$$

③ 8, 0.8, 0.08, 0.008, ...

$$a_1 = 8, r = \frac{1}{10}, \Rightarrow a_n = 8 \cdot \left(\frac{1}{10}\right)^{n-1}$$

④ 48, 24, 12, 6, 3, $\frac{3}{2}$, $\frac{3}{4}$, ...

$$a_1 = 48, r = \frac{1}{2}, \Rightarrow a_n = 48 \cdot \left(\frac{1}{2}\right)^{n-1}$$

Find the stated term of the given geometric sequence.

① 14th term of 5, -10, 20, -40, 80, ...

② 17th term of 3, 6, 12, 24, 48, ...

③ 12th term of 54, 18, 6, 2, ...

④ 10th term of 24, 36, 54, 81, ...

⑤ 7th term of 75, -30, 12, -4.8, ...

Geometric sequence

$$a_n = a_1 \cdot r^{n-1}$$

Find the closed form formula for the given geometric sequence.

① 7, 14, 28, 56, ...

$$a_1 = 7, r = 2, \Rightarrow a_n = 7 \cdot 2^{n-1}$$

② 4, -20, 100, -500, ...

$$a_1 = 4, r = -5, \Rightarrow a_n = 4 \cdot (-5)^{n-1}$$

③ 8, 0.8, 0.08, 0.008, ...

$$a_1 = 8, r = \frac{1}{10}, \Rightarrow a_n = 8 \cdot \left(\frac{1}{10}\right)^{n-1}$$

④ 48, 24, 12, 6, 3, $\frac{3}{2}$, $\frac{3}{4}$, ...

$$a_1 = 48, r = \frac{1}{2}, \Rightarrow a_n = 48 \cdot \left(\frac{1}{2}\right)^{n-1}$$

Find the stated term of the given geometric sequence.

① 14th term of 5, -10, 20, -40, 80, ...

$$a_n = 5 \cdot (-2)^{n-1}$$

$$\Rightarrow a_{14} = 5 \cdot (-2)^{13} = 5 \cdot (-8192) = -40,960$$

② 17th term of 3, 6, 12, 24, 48, ...

③ 12th term of 54, 18, 6, 2, ...

④ 10th term of 24, 36, 54, 81, ...

⑤ 7th term of 75, -30, 12, -4.8, ...

Geometric sequence

$$a_n = a_1 \cdot r^{n-1}$$

Find the closed form formula for the given geometric sequence.

① 7, 14, 28, 56, ...

$$a_1 = 7, r = 2, \Rightarrow a_n = 7 \cdot 2^{n-1}$$

② 4, -20, 100, -500, ...

$$a_1 = 4, r = -5, \Rightarrow a_n = 4 \cdot (-5)^{n-1}$$

③ 8, 0.8, 0.08, 0.008, ...

$$a_1 = 8, r = \frac{1}{10}, \Rightarrow a_n = 8 \cdot \left(\frac{1}{10}\right)^{n-1}$$

④ 48, 24, 12, 6, 3, $\frac{3}{2}$, $\frac{3}{4}$, ...

$$a_1 = 48, r = \frac{1}{2}, \Rightarrow a_n = 48 \cdot \left(\frac{1}{2}\right)^{n-1}$$

Find the stated term of the given geometric sequence.

① 14th term of 5, -10, 20, -40, 80, ...

$$a_n = 5 \cdot (-2)^{n-1}$$

$$\Rightarrow a_{14} = 5 \cdot (-2)^{13} = 5 \cdot (-8192) = -40,960$$

② 17th term of 3, 6, 12, 24, 48, ...

$$a_n = 3 \cdot 2^{n-1}$$

$$\Rightarrow a_{17} = 3 \cdot 2^{16} = 196,608$$

③ 12th term of 54, 18, 6, 2, ...

④ 10th term of 24, 36, 54, 81, ...

⑤ 7th term of 75, -30, 12, -4.8, ...

Geometric sequence

$$a_n = a_1 \cdot r^{n-1}$$

Find the closed form formula for the given geometric sequence.

- 7, 14, 28, 56, ...
 $a_1 = 7, r = 2, \Rightarrow a_n = 7 \cdot 2^{n-1}$
- 4, -20, 100, -500, ...
 $a_1 = 4, r = -5, \Rightarrow a_n = 4 \cdot (-5)^{n-1}$
- 8, 0.8, 0.08, 0.008, ...
 $a_1 = 8, r = \frac{1}{10}, \Rightarrow a_n = 8 \cdot \left(\frac{1}{10}\right)^{n-1}$
- 48, 24, 12, 6, 3, $\frac{3}{2}, \frac{3}{4}, \dots$
 $a_1 = 48, r = \frac{1}{2}, \Rightarrow a_n = 48 \cdot \left(\frac{1}{2}\right)^{n-1}$

Find the stated term of the given geometric sequence.

- 14th term of 5, -10, 20, -40, 80, ...
 $a_n = 5 \cdot (-2)^{n-1}$
 $\Rightarrow a_{14} = 5 \cdot (-2)^{13} = 5 \cdot (-8192) = -40,960$
- 17th term of 3, 6, 12, 24, 48, ...
 $a_n = 3 \cdot 2^{n-1}$
 $\Rightarrow a_{17} = 3 \cdot 2^{16} = 196,608$
- 12th term of 54, 18, 6, 2, ...
 $a_n = 54 \cdot \left(\frac{1}{3}\right)^{n-1}$
 $\Rightarrow a_{12} = 54 \cdot \left(\frac{1}{3}\right)^{11} = \frac{2 \cdot 27}{3^{11}} = \frac{2 \cdot 3^3}{3^{11}} = \frac{2}{3^8} = \frac{2}{6561}$
- 10th term of 24, 36, 54, 81, ...
 $a_n = 24 \cdot \left(\frac{3}{2}\right)^{n-1}$
 $\Rightarrow a_{10} = 24 \cdot \left(\frac{3}{2}\right)^9 = 24 \cdot \frac{3^9}{2^9} = 24 \cdot \frac{19683}{512} = \frac{472392}{128} = 3689.78125$
- 7th term of 75, -30, 12, -4.8, ...
 $a_n = 75 \cdot \left(-\frac{2}{5}\right)^{n-1}$
 $\Rightarrow a_7 = 75 \cdot \left(-\frac{2}{5}\right)^6 = 75 \cdot \frac{64}{15625} = \frac{4800}{15625} = \frac{96}{3125} = 0.03072$

Geometric sequence

$$a_n = a_1 \cdot r^{n-1}$$

Find the closed form formula for the given geometric sequence.

- ① 7, 14, 28, 56, ...
 $a_1 = 7, r = 2, \Rightarrow a_n = 7 \cdot 2^{n-1}$
- ② 4, -20, 100, -500, ...
 $a_1 = 4, r = -5, \Rightarrow a_n = 4 \cdot (-5)^{n-1}$
- ③ 8, 0.8, 0.08, 0.008, ...
 $a_1 = 8, r = \frac{1}{10}, \Rightarrow a_n = 8 \cdot \left(\frac{1}{10}\right)^{n-1}$
- ④ 48, 24, 12, 6, 3, $\frac{3}{2}, \frac{3}{4}, \dots$
 $a_1 = 48, r = \frac{1}{2}, \Rightarrow a_n = 48 \cdot \left(\frac{1}{2}\right)^{n-1}$

Find the stated term of the given geometric sequence.

- ① 14th term of 5, -10, 20, -40, 80, ...
 $a_n = 5 \cdot (-2)^{n-1}$
 $\Rightarrow a_{14} = 5 \cdot (-2)^{13} = 5 \cdot (-8192) = -40,960$
- ② 17th term of 3, 6, 12, 24, 48, ...
 $a_n = 3 \cdot 2^{n-1}$
 $\Rightarrow a_{17} = 3 \cdot 2^{16} = 196,608$
- ③ 12th term of 54, 18, 6, 2, ...
 $a_n = 54 \cdot \left(\frac{1}{3}\right)^{n-1}$
 $\Rightarrow a_{12} = 54 \cdot \left(\frac{1}{3}\right)^{11} = \frac{2 \cdot 27}{3^{11}} = \frac{2 \cdot 3^3}{3^{11}} = \frac{2}{3^8} = \frac{2}{6561}$
- ④ 10th term of 24, 36, 54, 81, ...
 $r = \frac{36}{24} = \frac{3}{2}, \Rightarrow a_n = 24 \cdot \left(\frac{3}{2}\right)^{n-1}$
 $\Rightarrow a_{10} = 24 \cdot \left(\frac{3}{2}\right)^9 = \frac{24 \cdot 3^9}{2^9} = \frac{472,392}{512} = \frac{59,049}{64}$
- ⑤ 7th term of 75, -30, 12, -4.8, ...

Geometric sequence

$$a_n = a_1 \cdot r^{n-1}$$

Find the closed form formula for the given geometric sequence.

- 7, 14, 28, 56, ...
 $a_1 = 7, r = 2, \Rightarrow a_n = 7 \cdot 2^{n-1}$
- 4, -20, 100, -500, ...
 $a_1 = 4, r = -5, \Rightarrow a_n = 4 \cdot (-5)^{n-1}$
- 8, 0.8, 0.08, 0.008, ...
 $a_1 = 8, r = \frac{1}{10}, \Rightarrow a_n = 8 \cdot \left(\frac{1}{10}\right)^{n-1}$
- 48, 24, 12, 6, 3, $\frac{3}{2}, \frac{3}{4}, \dots$
 $a_1 = 48, r = \frac{1}{2}, \Rightarrow a_n = 48 \cdot \left(\frac{1}{2}\right)^{n-1}$

Find the stated term of the given geometric sequence.

- 14th term of 5, -10, 20, -40, 80, ...
 $a_n = 5 \cdot (-2)^{n-1}$
 $\Rightarrow a_{14} = 5 \cdot (-2)^{13} = 5 \cdot (-8192) = -40,960$
- 17th term of 3, 6, 12, 24, 48, ...
 $a_n = 3 \cdot 2^{n-1}$
 $\Rightarrow a_{17} = 3 \cdot 2^{16} = 196,608$
- 12th term of 54, 18, 6, 2, ...
 $a_n = 54 \cdot \left(\frac{1}{3}\right)^{n-1}$
 $\Rightarrow a_{12} = 54 \cdot \left(\frac{1}{3}\right)^{11} = \frac{2 \cdot 27}{3^{11}} = \frac{2 \cdot 3^3}{3^{11}} = \frac{2}{3^8} = \frac{2}{6561}$
- 10th term of 24, 36, 54, 81, ...
 $r = \frac{36}{24} = \frac{3}{2}, \Rightarrow a_n = 24 \cdot \left(\frac{3}{2}\right)^{n-1}$
 $\Rightarrow a_{10} = 24 \cdot \left(\frac{3}{2}\right)^9 = \frac{24 \cdot 3^9}{2^9} = \frac{472,392}{512} = \frac{59,049}{64}$
- 7th term of 75, -30, 12, -4.8, ...
 $r = \frac{-30}{75} = -\frac{2}{5}, \Rightarrow a_n = 75 \cdot \left(-\frac{2}{5}\right)^{n-1}$
 $\Rightarrow a_7 = 75 \cdot \left(-\frac{2}{5}\right)^6 = \frac{75 \cdot 2^6}{5^6} = \frac{4,800}{15,625} = \frac{192}{625}$

Example

Find the sum of the first 4 terms of $3, 30, 300, 3000, 30000, 300000, \dots$

Example

Find the sum of the first 4 terms of $3, 30, 300, 3000, 30000, 300000, \dots$

Answer: Multiply by $(1 - 10)$ and use the “telescopic sum” trick:

$$(1 - 10) \cdot (3 + 30 + 300 + 3000)$$

Example

Find the sum of the first 4 terms of 3, 30, 300, 3000, 30000, 300000,

Answer: Multiply by $(1 - 10)$ and use the “telescopic sum” trick:

$$\begin{aligned}(1 - 10) \cdot (3 + 30 + 300 + 3000) \\ = 3 - 30 + 30 - 300 + 300 - 3000 + 3000 - 30000\end{aligned}$$

Example

Find the sum of the first 4 terms of 3, 30, 300, 3000, 30000, 300000, . . .

Answer: Multiply by $(1 - 10)$ and use the “telescopic sum” trick:

$$\begin{aligned} &(1 - 10) \cdot (3 + 30 + 300 + 3000) \\ &= 3 - \underbrace{30 + 30} - \underbrace{300 + 300} - \underbrace{3000 + 3000} - 30000 \end{aligned}$$

Example

Find the sum of the first 4 terms of 3, 30, 300, 3000, 30000, 300000, ...

Answer: Multiply by $(1 - 10)$ and use the “telescopic sum” trick:

$$\begin{aligned}(1 - 10) \cdot (3 + 30 + 300 + 3000) \\&= 3 - \underbrace{30 + 30} - \underbrace{300 + 300} - \underbrace{3000 + 3000} - 30000 \\&= 3 - 30000\end{aligned}$$

Example

Find the sum of the first 4 terms of 3, 30, 300, 3000, 30000, 300000, ...

Answer: Multiply by $(1 - 10)$ and use the “telescopic sum” trick:

$$\begin{aligned}(1 - 10) \cdot (3 + 30 + 300 + 3000) \\&= 3 - \underbrace{30 + 30}_{300} - \underbrace{300 + 300}_{3000} - \underbrace{3000 + 3000}_{30000} - 30000 \\&= 3 - 30000\end{aligned}$$

Therefore: $(1 - 10) \cdot \sum_{n=1}^4 a_n = 3 - 30000$

Geometric series

Example

Find the sum of the first 4 terms of 3, 30, 300, 3000, 30000, 300000, ...

Answer: Multiply by $(1 - 10)$ and use the “telescopic sum” trick:

$$\begin{aligned}(1 - 10) \cdot (3 + 30 + 300 + 3000) \\&= 3 - \underbrace{30 + 30} - \underbrace{300 + 300} - \underbrace{3000 + 3000} - 30000 \\&= 3 - 30000\end{aligned}$$

Therefore: $(1 - 10) \cdot \sum_{n=1}^4 a_n = 3 - 30000 \quad \Rightarrow \quad \sum_{n=1}^4 a_n = \frac{3 - 30000}{1 - 10} = 3 \cdot \frac{1 - 10000}{1 - 10}.$

Geometric series

Example

Find the sum of the first 4 terms of 3, 30, 300, 3000, 30000, 300000, ...

Answer: Multiply by $(1 - 10)$ and use the “telescopic sum” trick:

$$\begin{aligned}(1 - 10) \cdot (3 + 30 + 300 + 3000) \\&= 3 - \underbrace{30 + 30} - \underbrace{300 + 300} - \underbrace{3000 + 3000} - 30000 \\&= 3 - 30000\end{aligned}$$

Therefore: $(1 - 10) \cdot \sum_{n=1}^4 a_n = 3 - 30000 \quad \Rightarrow \quad \sum_{n=1}^4 a_n = \frac{3 - 30000}{1 - 10} = 3 \cdot \frac{1 - 10000}{1 - 10}.$

Evaluating this gives the expected answer: $\sum_{n=1}^4 a_n = 3 \cdot \frac{-9999}{-9} = 3 \cdot 1111 = 3333.$

Geometric series

Example

Find the sum of the first 4 terms of 3, 30, 300, 3000, 30000, 300000, ...

Answer: Multiply by $(1 - 10)$ and use the “telescopic sum” trick:

$$\begin{aligned}(1 - 10) \cdot (3 + 30 + 300 + 3000) \\&= 3 - \underbrace{30 + 30} - \underbrace{300 + 300} - \underbrace{3000 + 3000} - 30000 \\&= 3 - 30000\end{aligned}$$

Therefore: $(1 - 10) \cdot \sum_{n=1}^4 a_n = 3 - 30000 \quad \Rightarrow \quad \sum_{n=1}^4 a_n = \frac{3 - 30000}{1 - 10} = 3 \cdot \frac{1 - 10000}{1 - 10}.$

Evaluating this gives the expected answer: $\sum_{n=1}^4 a_n = 3 \cdot \frac{-9999}{-9} = 3 \cdot 1111 = 3333.$

Geometric series

For any geometric series $a_n = a_1 \cdot r^{n-1}$ the sum of the first p terms is

$$\sum_{n=1}^p a_n = a_1 \cdot \frac{1 - r^p}{1 - r}$$

Geometric series - exercises

Geometric series

For $a_n = a_1 \cdot r^{n-1}$:

$$\sum_{n=1}^p a_n = a_1 \cdot \frac{1 - r^p}{1 - r}$$

1 Find $\sum_{n=1}^9 4 \cdot 6^{n-1}$

2 Find $\sum_{n=1}^6 5 \cdot (-3)^{n-1}$

Find the sum of the given geometric sequence.

1 Sum the first 7 terms of $2, -10, 50, -250, \dots$

2 Sum the first 8 terms of $9, 6, 4, \frac{8}{3}, \dots$

3 Sum the first 15 terms of $7, -\frac{7}{3}, \frac{7}{9}, -\frac{7}{27}, \dots$

Geometric series - exercises

Geometric series

For $a_n = a_1 \cdot r^{n-1}$:

$$\sum_{n=1}^p a_n = a_1 \cdot \frac{1 - r^p}{1 - r}$$

① Find $\sum_{n=1}^9 4 \cdot 6^{n-1}$
 $p = 9, \quad a_1 = 4, \quad r = 6$

② Find $\sum_{n=1}^6 5 \cdot (-3)^{n-1}$

Find the sum of the given geometric sequence.

① Sum the first 7 terms of $2, -10, 50, -250, \dots$

② Sum the first 8 terms of $9, 6, 4, \frac{8}{3}, \dots$

③ Sum the first 15 terms of $7, -\frac{7}{3}, \frac{7}{9}, -\frac{7}{27}, \dots$

Geometric series - exercises

Geometric series

For $a_n = a_1 \cdot r^{n-1}$:

$$\sum_{n=1}^p a_n = a_1 \cdot \frac{1 - r^p}{1 - r}$$

1 Find $\sum_{n=1}^9 4 \cdot 6^{n-1}$

$$p = 9, \quad a_1 = 4, \quad r = 6$$

$$\begin{aligned} \Rightarrow \sum_{n=1}^9 4 \cdot 6^{n-1} &= 4 \cdot \frac{1-6^9}{1-6} \\ &= \frac{4 \cdot (-10,077,695)}{-5} = 8,062,156 \end{aligned}$$

2 Find $\sum_{n=1}^6 5 \cdot (-3)^{n-1}$

Find the sum of the given geometric sequence.

1 Sum the first 7 terms of $2, -10, 50, -250, \dots$

2 Sum the first 8 terms of $9, 6, 4, \frac{8}{3}, \dots$

3 Sum the first 15 terms of $7, -\frac{7}{3}, \frac{7}{9}, -\frac{7}{27}, \dots$

Geometric series - exercises

Geometric series

For $a_n = a_1 \cdot r^{n-1}$:

$$\sum_{n=1}^p a_n = a_1 \cdot \frac{1 - r^p}{1 - r}$$

① Find $\sum_{n=1}^9 4 \cdot 6^{n-1}$

$$p = 9, \quad a_1 = 4, \quad r = 6$$

$$\begin{aligned} \Rightarrow \sum_{n=1}^9 4 \cdot 6^{n-1} &= 4 \cdot \frac{1-6^9}{1-6} \\ &= \frac{4 \cdot (-10,077,695)}{-5} = 8,062,156 \end{aligned}$$

② Find $\sum_{n=1}^6 5 \cdot (-3)^{n-1}$

$$p = 6, \quad a_1 = 5, \quad r = -3$$

Find the sum of the given geometric sequence.

① Sum the first 7 terms of $2, -10, 50, -250, \dots$

② Sum the first 8 terms of $9, 6, 4, \frac{8}{3}, \dots$

③ Sum the first 15 terms of $7, -\frac{7}{3}, \frac{7}{9}, -\frac{7}{27}, \dots$

Geometric series - exercises

Geometric series

For $a_n = a_1 \cdot r^{n-1}$:

$$\sum_{n=1}^p a_n = a_1 \cdot \frac{1 - r^p}{1 - r}$$

① Find $\sum_{n=1}^9 4 \cdot 6^{n-1}$

$$p = 9, \quad a_1 = 4, \quad r = 6$$

$$\begin{aligned} \Rightarrow \sum_{n=1}^9 4 \cdot 6^{n-1} &= 4 \cdot \frac{1-6^9}{1-6} \\ &= \frac{4 \cdot (-10,077,695)}{-5} = 8,062,156 \end{aligned}$$

② Find $\sum_{n=1}^6 5 \cdot (-3)^{n-1}$

$$p = 6, \quad a_1 = 5, \quad r = -3$$

$$\begin{aligned} \Rightarrow \sum_{n=1}^6 5 \cdot (-3)^{n-1} &= 5 \cdot \frac{1-(-3)^6}{1-(-3)} = 5 \cdot \frac{1-729}{1+3} \\ &= 5 \cdot \frac{-728}{4} = -910 \end{aligned}$$

Find the sum of the given geometric sequence.

① Sum the first 7 terms of 2, -10, 50, -250, ...

② Sum the first 8 terms of 9, 6, 4, $\frac{8}{3}$, ...

③ Sum the first 15 terms of 7, $-\frac{7}{3}$, $\frac{7}{9}$, $-\frac{7}{27}$, ...

Geometric series - exercises

Geometric series

For $a_n = a_1 \cdot r^{n-1}$:

$$\sum_{n=1}^p a_n = a_1 \cdot \frac{1 - r^p}{1 - r}$$

① Find $\sum_{n=1}^9 4 \cdot 6^{n-1}$

$$p = 9, \quad a_1 = 4, \quad r = 6$$

$$\begin{aligned} \Rightarrow \sum_{n=1}^9 4 \cdot 6^{n-1} &= 4 \cdot \frac{1-6^9}{1-6} \\ &= \frac{4 \cdot (-10,077,695)}{-5} = 8,062,156 \end{aligned}$$

② Find $\sum_{n=1}^6 5 \cdot (-3)^{n-1}$

$$p = 6, \quad a_1 = 5, \quad r = -3$$

$$\begin{aligned} \Rightarrow \sum_{n=1}^6 5 \cdot (-3)^{n-1} &= 5 \cdot \frac{1-(-3)^6}{1-(-3)} = 5 \cdot \frac{1-729}{1+3} \\ &= 5 \cdot \frac{-728}{4} = -910 \end{aligned}$$

Find the sum of the given geometric sequence.

① Sum the first 7 terms of 2, -10, 50, -250, ...

We need $\sum_{n=1}^7 a_n$ for $a_n = 2 \cdot (-5)^{n-1}$.

② Sum the first 8 terms of 9, 6, 4, $\frac{8}{3}$, ...

③ Sum the first 15 terms of 7, $-\frac{7}{3}$, $\frac{7}{9}$, $-\frac{7}{27}$, ...

Geometric series - exercises

Geometric series

For $a_n = a_1 \cdot r^{n-1}$:

$$\sum_{n=1}^p a_n = a_1 \cdot \frac{1 - r^p}{1 - r}$$

① Find $\sum_{n=1}^9 4 \cdot 6^{n-1}$

$$p = 9, \quad a_1 = 4, \quad r = 6$$

$$\begin{aligned} \Rightarrow \sum_{n=1}^9 4 \cdot 6^{n-1} &= 4 \cdot \frac{1-6^9}{1-6} \\ &= \frac{4 \cdot (-10,077,695)}{-5} = 8,062,156 \end{aligned}$$

② Find $\sum_{n=1}^6 5 \cdot (-3)^{n-1}$

$$p = 6, \quad a_1 = 5, \quad r = -3$$

$$\begin{aligned} \Rightarrow \sum_{n=1}^6 5 \cdot (-3)^{n-1} &= 5 \cdot \frac{1-(-3)^6}{1-(-3)} = 5 \cdot \frac{1-729}{1+3} \\ &= 5 \cdot \frac{-728}{4} = -910 \end{aligned}$$

Find the sum of the given geometric sequence.

① Sum the first 7 terms of 2, -10, 50, -250, ...

We need $\sum_{n=1}^7 a_n$ for $a_n = 2 \cdot (-5)^{n-1}$.

$$p = 7, \quad a_1 = 2, \quad r = -5$$

② Sum the first 8 terms of 9, 6, 4, $\frac{8}{3}$, ...

③ Sum the first 15 terms of 7, $-\frac{7}{3}$, $\frac{7}{9}$, $-\frac{7}{27}$, ...

Geometric series - exercises

Geometric series

For $a_n = a_1 \cdot r^{n-1}$:

$$\sum_{n=1}^p a_n = a_1 \cdot \frac{1 - r^p}{1 - r}$$

① Find $\sum_{n=1}^9 4 \cdot 6^{n-1}$

$$p = 9, \quad a_1 = 4, \quad r = 6$$

$$\begin{aligned} \Rightarrow \sum_{n=1}^9 4 \cdot 6^{n-1} &= 4 \cdot \frac{1-6^9}{1-6} \\ &= \frac{4 \cdot (-10,077,695)}{-5} = 8,062,156 \end{aligned}$$

② Find $\sum_{n=1}^6 5 \cdot (-3)^{n-1}$

$$p = 6, \quad a_1 = 5, \quad r = -3$$

$$\begin{aligned} \Rightarrow \sum_{n=1}^6 5 \cdot (-3)^{n-1} &= 5 \cdot \frac{1-(-3)^6}{1-(-3)} = 5 \cdot \frac{1-729}{1+3} \\ &= 5 \cdot \frac{-728}{4} = -910 \end{aligned}$$

Find the sum of the given geometric sequence.

① Sum the first 7 terms of 2, -10, 50, -250, ...

We need $\sum_{n=1}^7 a_n$ for $a_n = 2 \cdot (-5)^{n-1}$.

$$p = 7, \quad a_1 = 2, \quad r = -5$$

$$\Rightarrow \sum_{n=1}^7 a_n = 2 \cdot \frac{1-(-5)^7}{1-(-5)} = 2 \cdot \frac{1+78125}{1+5} = 26,042$$

② Sum the first 8 terms of 9, 6, 4, $\frac{8}{3}$, ...

③ Sum the first 15 terms of 7, $-\frac{7}{3}$, $\frac{7}{9}$, $-\frac{7}{27}$, ...

Geometric series - exercises

Geometric series

For $a_n = a_1 \cdot r^{n-1}$:

$$\sum_{n=1}^p a_n = a_1 \cdot \frac{1 - r^p}{1 - r}$$

① Find $\sum_{n=1}^9 4 \cdot 6^{n-1}$

$$p = 9, \quad a_1 = 4, \quad r = 6$$

$$\begin{aligned} \Rightarrow \sum_{n=1}^9 4 \cdot 6^{n-1} &= 4 \cdot \frac{1-6^9}{1-6} \\ &= \frac{4 \cdot (-10,077,695)}{-5} = 8,062,156 \end{aligned}$$

② Find $\sum_{n=1}^6 5 \cdot (-3)^{n-1}$

$$p = 6, \quad a_1 = 5, \quad r = -3$$

$$\begin{aligned} \Rightarrow \sum_{n=1}^6 5 \cdot (-3)^{n-1} &= 5 \cdot \frac{1-(-3)^6}{1-(-3)} = 5 \cdot \frac{1-729}{1+3} \\ &= 5 \cdot \frac{-728}{4} = -910 \end{aligned}$$

Find the sum of the given geometric sequence.

① Sum the first 7 terms of $2, -10, 50, -250, \dots$

We need $\sum_{n=1}^7 a_n$ for $a_n = 2 \cdot (-5)^{n-1}$.

$$p = 7, \quad a_1 = 2, \quad r = -5$$

$$\Rightarrow \sum_{n=1}^7 a_n = 2 \cdot \frac{1-(-5)^7}{1-(-5)} = 2 \cdot \frac{1+78125}{1+5} = 26,042$$

② Sum the first 8 terms of $9, 6, 4, \frac{8}{3}, \dots$

We need $\sum_{n=1}^8 a_n$ for $a_n = 9 \cdot \left(\frac{2}{3}\right)^{n-1}$.

③ Sum the first 15 terms of $7, -\frac{7}{3}, \frac{7}{9}, -\frac{7}{27}, \dots$

Geometric series - exercises

Geometric series

For $a_n = a_1 \cdot r^{n-1}$:

$$\sum_{n=1}^p a_n = a_1 \cdot \frac{1 - r^p}{1 - r}$$

① Find $\sum_{n=1}^9 4 \cdot 6^{n-1}$

$$p = 9, \quad a_1 = 4, \quad r = 6$$

$$\begin{aligned} \Rightarrow \sum_{n=1}^9 4 \cdot 6^{n-1} &= 4 \cdot \frac{1-6^9}{1-6} \\ &= \frac{4 \cdot (-10,077,695)}{-5} = 8,062,156 \end{aligned}$$

② Find $\sum_{n=1}^6 5 \cdot (-3)^{n-1}$

$$p = 6, \quad a_1 = 5, \quad r = -3$$

$$\begin{aligned} \Rightarrow \sum_{n=1}^6 5 \cdot (-3)^{n-1} &= 5 \cdot \frac{1-(-3)^6}{1-(-3)} = 5 \cdot \frac{1-729}{1+3} \\ &= 5 \cdot \frac{-728}{4} = -910 \end{aligned}$$

Find the sum of the given geometric sequence.

① Sum the first 7 terms of $2, -10, 50, -250, \dots$

We need $\sum_{n=1}^7 a_n$ for $a_n = 2 \cdot (-5)^{n-1}$.

$$p = 7, \quad a_1 = 2, \quad r = -5$$

$$\Rightarrow \sum_{n=1}^7 a_n = 2 \cdot \frac{1-(-5)^7}{1-(-5)} = 2 \cdot \frac{1+78125}{1+5} = 26,042$$

② Sum the first 8 terms of $9, 6, 4, \frac{8}{3}, \dots$

We need $\sum_{n=1}^8 a_n$ for $a_n = 9 \cdot (\frac{2}{3})^{n-1}$.

$$p = 8, \quad a_1 = 9, \quad r = \frac{2}{3}$$

③ Sum the first 15 terms of $7, -\frac{7}{3}, \frac{7}{9}, -\frac{7}{27}, \dots$

Geometric series - exercises

Geometric series

For $a_n = a_1 \cdot r^{n-1}$:

$$\sum_{n=1}^p a_n = a_1 \cdot \frac{1 - r^p}{1 - r}$$

① Find $\sum_{n=1}^9 4 \cdot 6^{n-1}$

$$p = 9, \quad a_1 = 4, \quad r = 6$$

$$\begin{aligned} \Rightarrow \sum_{n=1}^9 4 \cdot 6^{n-1} &= 4 \cdot \frac{1-6^9}{1-6} \\ &= \frac{4 \cdot (-10,077,695)}{-5} = 8,062,156 \end{aligned}$$

② Find $\sum_{n=1}^6 5 \cdot (-3)^{n-1}$

$$p = 6, \quad a_1 = 5, \quad r = -3$$

$$\begin{aligned} \Rightarrow \sum_{n=1}^6 5 \cdot (-3)^{n-1} \\ &= 5 \cdot \frac{1-(-3)^6}{1-(-3)} = 5 \cdot \frac{1-729}{1+3} \\ &= 5 \cdot \frac{-728}{4} = -910 \end{aligned}$$

Find the sum of the given geometric sequence.

① Sum the first 7 terms of $2, -10, 50, -250, \dots$

We need $\sum_{n=1}^7 a_n$ for $a_n = 2 \cdot (-5)^{n-1}$.

$$p = 7, \quad a_1 = 2, \quad r = -5$$

$$\Rightarrow \sum_{n=1}^7 a_n = 2 \cdot \frac{1-(-5)^7}{1-(-5)} = 2 \cdot \frac{1+78125}{1+5} = 26,042$$

② Sum the first 8 terms of $9, 6, 4, \frac{8}{3}, \dots$

We need $\sum_{n=1}^8 a_n$ for $a_n = 9 \cdot (\frac{2}{3})^{n-1}$.

$$p = 8, \quad a_1 = 9, \quad r = \frac{2}{3}$$

$$\begin{aligned} \Rightarrow \sum_{n=1}^8 a_n &= 9 \cdot \frac{1-(\frac{2}{3})^8}{1-\frac{2}{3}} = \frac{9 \cdot (1-\frac{256}{6561})}{\frac{1}{3}} \\ &= 9 \cdot \frac{6561-256}{6561} \cdot \frac{3}{1} = \frac{9 \cdot 6205 \cdot 3}{6561} = \frac{6205}{243} \end{aligned}$$

③ Sum the first 15 terms of $7, -\frac{7}{3}, \frac{7}{9}, -\frac{7}{27}, \dots$

Geometric series - exercises

Geometric series

For $a_n = a_1 \cdot r^{n-1}$:

$$\sum_{n=1}^p a_n = a_1 \cdot \frac{1 - r^p}{1 - r}$$

① Find $\sum_{n=1}^9 4 \cdot 6^{n-1}$

$$p = 9, \quad a_1 = 4, \quad r = 6$$

$$\begin{aligned} \Rightarrow \sum_{n=1}^9 4 \cdot 6^{n-1} &= 4 \cdot \frac{1-6^9}{1-6} \\ &= \frac{4 \cdot (-10,077,695)}{-5} = 8,062,156 \end{aligned}$$

② Find $\sum_{n=1}^6 5 \cdot (-3)^{n-1}$

$$p = 6, \quad a_1 = 5, \quad r = -3$$

$$\begin{aligned} \Rightarrow \sum_{n=1}^6 5 \cdot (-3)^{n-1} \\ &= 5 \cdot \frac{1-(-3)^6}{1-(-3)} = 5 \cdot \frac{1-729}{1+3} \\ &= 5 \cdot \frac{-728}{4} = -910 \end{aligned}$$

Find the sum of the given geometric sequence.

① Sum the first 7 terms of 2, -10, 50, -250, ...

We need $\sum_{n=1}^7 a_n$ for $a_n = 2 \cdot (-5)^{n-1}$.

$$p = 7, \quad a_1 = 2, \quad r = -5$$

$$\Rightarrow \sum_{n=1}^7 a_n = 2 \cdot \frac{1-(-5)^7}{1-(-5)} = 2 \cdot \frac{1+78125}{1+5} = 26,042$$

② Sum the first 8 terms of 9, 6, 4, $\frac{8}{3}$, ...

We need $\sum_{n=1}^8 a_n$ for $a_n = 9 \cdot (\frac{2}{3})^{n-1}$.

$$p = 8, \quad a_1 = 9, \quad r = \frac{2}{3}$$

$$\begin{aligned} \Rightarrow \sum_{n=1}^8 a_n &= 9 \cdot \frac{1-(\frac{2}{3})^8}{1-\frac{2}{3}} = \frac{9 \cdot (1-\frac{256}{6561})}{\frac{1}{3}} \\ &= 9 \cdot \frac{6561-256}{6561} \cdot \frac{3}{1} = \frac{9 \cdot 6205 \cdot 3}{6561} = \frac{6205}{243} \end{aligned}$$

③ Sum the first 15 terms of 7, $-\frac{7}{3}$, $\frac{7}{9}$, $-\frac{7}{27}$, ...

We need $\sum_{n=1}^{15} a_n$ for $a_n = 7 \cdot (-\frac{1}{3})^{n-1}$.

Geometric series - exercises

Geometric series

For $a_n = a_1 \cdot r^{n-1}$:

$$\sum_{n=1}^p a_n = a_1 \cdot \frac{1 - r^p}{1 - r}$$

① Find $\sum_{n=1}^9 4 \cdot 6^{n-1}$

$$p = 9, \quad a_1 = 4, \quad r = 6$$

$$\begin{aligned} \Rightarrow \sum_{n=1}^9 4 \cdot 6^{n-1} &= 4 \cdot \frac{1-6^9}{1-6} \\ &= \frac{4 \cdot (-10,077,695)}{-5} = 8,062,156 \end{aligned}$$

② Find $\sum_{n=1}^6 5 \cdot (-3)^{n-1}$

$$p = 6, \quad a_1 = 5, \quad r = -3$$

$$\begin{aligned} \Rightarrow \sum_{n=1}^6 5 \cdot (-3)^{n-1} \\ &= 5 \cdot \frac{1-(-3)^6}{1-(-3)} = 5 \cdot \frac{1-729}{1+3} \\ &= 5 \cdot \frac{-728}{4} = -910 \end{aligned}$$

Find the sum of the given geometric sequence.

① Sum the first 7 terms of 2, -10, 50, -250, ...

We need $\sum_{n=1}^7 a_n$ for $a_n = 2 \cdot (-5)^{n-1}$.

$$p = 7, \quad a_1 = 2, \quad r = -5$$

$$\Rightarrow \sum_{n=1}^7 a_n = 2 \cdot \frac{1-(-5)^7}{1-(-5)} = 2 \cdot \frac{1+78125}{1+5} = 26,042$$

② Sum the first 8 terms of 9, 6, 4, $\frac{8}{3}$, ...

We need $\sum_{n=1}^8 a_n$ for $a_n = 9 \cdot (\frac{2}{3})^{n-1}$.

$$p = 8, \quad a_1 = 9, \quad r = \frac{2}{3}$$

$$\begin{aligned} \Rightarrow \sum_{n=1}^8 a_n &= 9 \cdot \frac{1-(\frac{2}{3})^8}{1-\frac{2}{3}} = \frac{9 \cdot (1-\frac{256}{6561})}{\frac{1}{3}} \\ &= 9 \cdot \frac{6561-256}{6561} \cdot \frac{3}{1} = \frac{9 \cdot 6205 \cdot 3}{6561} = \frac{6205}{243} \end{aligned}$$

③ Sum the first 15 terms of 7, $-\frac{7}{3}$, $\frac{7}{9}$, $-\frac{7}{27}$, ...

We need $\sum_{n=1}^{15} a_n$ for $a_n = 7 \cdot (-\frac{1}{3})^{n-1}$.

$$p = 15, \quad a_1 = 7, \quad r = -\frac{1}{3}$$

Geometric series - exercises

Geometric series

For $a_n = a_1 \cdot r^{n-1}$:

$$\sum_{n=1}^p a_n = a_1 \cdot \frac{1 - r^p}{1 - r}$$

① Find $\sum_{n=1}^9 4 \cdot 6^{n-1}$

$$p = 9, \quad a_1 = 4, \quad r = 6$$

$$\begin{aligned} \Rightarrow \sum_{n=1}^9 4 \cdot 6^{n-1} &= 4 \cdot \frac{1-6^9}{1-6} \\ &= \frac{4 \cdot (-10,077,695)}{-5} = 8,062,156 \end{aligned}$$

② Find $\sum_{n=1}^6 5 \cdot (-3)^{n-1}$

$$p = 6, \quad a_1 = 5, \quad r = -3$$

$$\begin{aligned} \Rightarrow \sum_{n=1}^6 5 \cdot (-3)^{n-1} \\ &= 5 \cdot \frac{1-(-3)^6}{1-(-3)} = 5 \cdot \frac{1-729}{1+3} \\ &= 5 \cdot \frac{-728}{4} = -910 \end{aligned}$$

Find the sum of the given geometric sequence.

① Sum the first 7 terms of 2, -10, 50, -250, ...

We need $\sum_{n=1}^7 a_n$ for $a_n = 2 \cdot (-5)^{n-1}$.

$$p = 7, \quad a_1 = 2, \quad r = -5$$

$$\Rightarrow \sum_{n=1}^7 a_n = 2 \cdot \frac{1-(-5)^7}{1-(-5)} = 2 \cdot \frac{1+78125}{1+5} = 26,042$$

② Sum the first 8 terms of 9, 6, 4, $\frac{8}{3}$, ...

We need $\sum_{n=1}^8 a_n$ for $a_n = 9 \cdot (\frac{2}{3})^{n-1}$.

$$p = 8, \quad a_1 = 9, \quad r = \frac{2}{3}$$

$$\begin{aligned} \Rightarrow \sum_{n=1}^8 a_n &= 9 \cdot \frac{1-(\frac{2}{3})^8}{1-\frac{2}{3}} = \frac{9 \cdot (1-\frac{256}{6561})}{\frac{1}{3}} \\ &= 9 \cdot \frac{6561-256}{6561} \cdot \frac{3}{1} = \frac{9 \cdot 6205 \cdot 3}{6561} = \frac{6205}{243} \end{aligned}$$

③ Sum the first 15 terms of 7, $-\frac{7}{3}$, $\frac{7}{9}$, $-\frac{7}{27}$, ...

We need $\sum_{n=1}^{15} a_n$ for $a_n = 7 \cdot (-\frac{1}{3})^{n-1}$.

$$p = 15, \quad a_1 = 7, \quad r = -\frac{1}{3}$$

$$\begin{aligned} \Rightarrow \sum_{n=1}^{15} a_n &= 7 \cdot \frac{1-(-\frac{1}{3})^{15}}{1-(-\frac{1}{3})} = \frac{7(1+\frac{1}{3^{15}})}{\frac{4}{3}} = 7 \cdot \frac{3^{15}+1}{3^{15}} \cdot \frac{3}{4} \\ &= \frac{7 \cdot 14,348,908}{3^{14} \cdot 4} = \frac{7 \cdot 3,587,227}{3^{14}} = \frac{25,110,589}{4,782,969} \end{aligned}$$

Infinite geometric series

Example

Add **all infinitely many** terms in the geometric series


$$\frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \frac{1}{16} + \frac{1}{32} + \dots$$

Infinite geometric series

Example

Add **all infinitely many** terms in the geometric series $\frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \frac{1}{16} + \frac{1}{32} + \dots$

Answer:




Infinite geometric series

Example

Add **all infinitely many** terms in the geometric series $\frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \frac{1}{16} + \frac{1}{32} + \dots$

Answer:




Infinite geometric series

Example

Add **all infinitely many** terms in the geometric series $\frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \frac{1}{16} + \frac{1}{32} + \dots$

Answer:




Infinite geometric series

Example

Add **all infinitely many** terms in the geometric series $\frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \frac{1}{16} + \frac{1}{32} + \dots$

Answer:




Infinite geometric series

Example

Add **all infinitely many** terms in the geometric series $\frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \frac{1}{16} + \frac{1}{32} + \dots$

Answer:




Infinite geometric series

Example

Add **all infinitely many** terms in the geometric series

$$\frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \frac{1}{16} + \frac{1}{32} + \dots$$

Answer:




Infinite geometric series

Example

Add **all infinitely many** terms in the geometric series

$$\frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \frac{1}{16} + \frac{1}{32} + \dots$$

Answer:




Infinite geometric series

Example

Add **all infinitely many** terms in the geometric series $\frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \frac{1}{16} + \frac{1}{32} + \dots$

Answer:




Infinite geometric series

Example

Add **all infinitely many** terms in the geometric series

$$\frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \frac{1}{16} + \frac{1}{32} + \dots$$

Answer:




Infinite geometric series

Example

Add **all infinitely many** terms in the geometric series $\frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \frac{1}{16} + \frac{1}{32} + \dots$

Answer:



Therefore:


$$\frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \frac{1}{16} + \frac{1}{32} + \dots = 1$$

Infinite geometric series

Example

Add **all infinitely many** terms in the geometric series $\frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \frac{1}{16} + \frac{1}{32} + \dots$

Answer:



Therefore:

$$\frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \frac{1}{16} + \frac{1}{32} + \dots = 1$$

Infinite geometric series

For any geometric sequence $a_1, a_1 \cdot r, a_1 \cdot r^2, a_1 \cdot r^3, \dots$
where the common ratio r is between $-1 < r < 1$, then the sum of all terms is

$$\sum_{n=1}^{\infty} a_n = a_1 \cdot \frac{1}{1-r}$$


Formal reason for this: $\sum_{n=1}^{\infty} a_n = \lim_{p \rightarrow \infty} \sum_{n=1}^p a_n = \lim_{p \rightarrow \infty} a_1 \cdot \frac{1-r^p}{1-r} = a_1 \cdot \frac{1}{1-r}$.

Infinite geometric series

Example

Add **all infinitely many** terms in the geometric series $\frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \frac{1}{16} + \frac{1}{32} + \dots$

Answer:



Therefore:

$$\frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \frac{1}{16} + \frac{1}{32} + \dots = 1$$

Infinite geometric series

For any geometric sequence $a_1, a_1 \cdot r, a_1 \cdot r^2, a_1 \cdot r^3, \dots$
where the common ratio r is between $-1 < r < 1$, then the sum of all terms is

$$\sum_{n=1}^{\infty} a_n = a_1 \cdot \frac{1}{1-r}$$

Formal reason for this: $\sum_{n=1}^{\infty} a_n = \lim_{p \rightarrow \infty} \sum_{n=1}^p a_n = \lim_{p \rightarrow \infty} a_1 \cdot \frac{1-r^p}{1-r} = a_1 \cdot \frac{1}{1-r}$.

Note: If the common ratio $r \geq 1$ or $r \leq -1$, then the infinite sum **does not exist!**

Example: The sum $5 + 10 + 20 + 40 + 80 + \dots$ does not exist!

Infinite geometric series - exercises

Infinite geometric series

For $a_1, a_1r, a_1r^2, a_1r^3, \dots$:

$$\sum_{n=1}^{\infty} a_n = a_1 \cdot \frac{1}{1-r}$$

1 Find $\sum_{n=1}^{\infty} 6 \cdot \left(\frac{1}{3}\right)^{n-1}$

2 Find $\sum_{n=1}^{\infty} 20 \cdot \left(\frac{2}{5}\right)^{n-1}$

3 Find $\sum_{n=1}^{\infty} 24 \cdot \left(-\frac{1}{2}\right)^{n-1}$

Find the sum of the given geometric sequence.

1 Sum all terms of $12, 3, \frac{3}{4}, \frac{3}{16}, \dots$

2 Sum all terms of $-27, 9, -3, 1, \dots$

3 Sum all terms of $50, -30, 18, \dots$

4 Sum all terms of $36, 24, 16, \dots$

5 Sum all terms of $0.4, 0.04, 0.004, 0.0004, \dots$

Infinite geometric series - exercises

Infinite geometric series

For $a_1, a_1r, a_1r^2, a_1r^3, \dots$:

$$\sum_{n=1}^{\infty} a_n = a_1 \cdot \frac{1}{1-r}$$

1 Find $\sum_{n=1}^{\infty} 6 \cdot \left(\frac{1}{3}\right)^{n-1}$
 $a_1 = 6, r = \frac{1}{3}$

2 Find $\sum_{n=1}^{\infty} 20 \cdot \left(\frac{2}{5}\right)^{n-1}$

3 Find $\sum_{n=1}^{\infty} 24 \cdot \left(-\frac{1}{2}\right)^{n-1}$

Find the sum of the given geometric sequence.

1 Sum all terms of $12, 3, \frac{3}{4}, \frac{3}{16}, \dots$

2 Sum all terms of $-27, 9, -3, 1, \dots$

3 Sum all terms of $50, -30, 18, \dots$

4 Sum all terms of $36, 24, 16, \dots$

5 Sum all terms of $0.4, 0.04, 0.004, 0.0004, \dots$

Infinite geometric series - exercises

Infinite geometric series

For $a_1, a_1r, a_1r^2, a_1r^3, \dots$:

$$\sum_{n=1}^{\infty} a_n = a_1 \cdot \frac{1}{1-r}$$

1 Find $\sum_{n=1}^{\infty} 6 \cdot \left(\frac{1}{3}\right)^{n-1}$

$$a_1 = 6, r = \frac{1}{3}$$

$$\Rightarrow \sum_{n=1}^{\infty} a_n = 6 \cdot \frac{1}{1-\frac{1}{3}} = \frac{6}{\frac{2}{3}} = 6 \cdot \frac{3}{2} = \frac{18}{2} = 9$$

2 Find $\sum_{n=1}^{\infty} 20 \cdot \left(\frac{2}{5}\right)^{n-1}$

3 Find $\sum_{n=1}^{\infty} 24 \cdot \left(-\frac{1}{2}\right)^{n-1}$

Find the sum of the given geometric sequence.

1 Sum all terms of $12, 3, \frac{3}{4}, \frac{3}{16}, \dots$

2 Sum all terms of $-27, 9, -3, 1, \dots$

3 Sum all terms of $50, -30, 18, \dots$

4 Sum all terms of $36, 24, 16, \dots$

5 Sum all terms of $0.4, 0.04, 0.004, 0.0004, \dots$

Infinite geometric series - exercises

Infinite geometric series

For $a_1, a_1r, a_1r^2, a_1r^3, \dots$:

$$\sum_{n=1}^{\infty} a_n = a_1 \cdot \frac{1}{1-r}$$

1 Find $\sum_{n=1}^{\infty} 6 \cdot \left(\frac{1}{3}\right)^{n-1}$

$$a_1 = 6, r = \frac{1}{3}$$

$$\Rightarrow \sum_{n=1}^{\infty} a_n = 6 \cdot \frac{1}{1-\frac{1}{3}} = \frac{6}{\frac{2}{3}} = 6 \cdot \frac{3}{2} = \frac{18}{2} = 9$$

2 Find $\sum_{n=1}^{\infty} 20 \cdot \left(\frac{2}{5}\right)^{n-1}$

$$a_1 = 20, r = \frac{2}{5}$$

3 Find $\sum_{n=1}^{\infty} 24 \cdot \left(-\frac{1}{2}\right)^{n-1}$

Find the sum of the given geometric sequence.

1 Sum all terms of $12, 3, \frac{3}{4}, \frac{3}{16}, \dots$

2 Sum all terms of $-27, 9, -3, 1, \dots$

3 Sum all terms of $50, -30, 18, \dots$

4 Sum all terms of $36, 24, 16, \dots$

5 Sum all terms of $0.4, 0.04, 0.004, 0.0004, \dots$

Infinite geometric series - exercises

Infinite geometric series

For $a_1, a_1r, a_1r^2, a_1r^3, \dots$:

$$\sum_{n=1}^{\infty} a_n = a_1 \cdot \frac{1}{1-r}$$

1 Find $\sum_{n=1}^{\infty} 6 \cdot \left(\frac{1}{3}\right)^{n-1}$

$$a_1 = 6, r = \frac{1}{3}$$

$$\Rightarrow \sum_{n=1}^{\infty} a_n = 6 \cdot \frac{1}{1-\frac{1}{3}} = \frac{6}{\frac{2}{3}} = 6 \cdot \frac{3}{2} = \frac{18}{2} = 9$$

2 Find $\sum_{n=1}^{\infty} 20 \cdot \left(\frac{2}{5}\right)^{n-1}$

$$a_1 = 20, r = \frac{2}{5}$$

$$\Rightarrow \sum_{n=1}^{\infty} a_n = 20 \cdot \frac{1}{1-\frac{2}{5}} = \frac{20}{\frac{3}{5}} = 20 \cdot \frac{5}{3} = \frac{100}{3}$$

3 Find $\sum_{n=1}^{\infty} 24 \cdot \left(-\frac{1}{2}\right)^{n-1}$

Find the sum of the given geometric sequence.

1 Sum all terms of $12, 3, \frac{3}{4}, \frac{3}{16}, \dots$

2 Sum all terms of $-27, 9, -3, 1, \dots$

3 Sum all terms of $50, -30, 18, \dots$

4 Sum all terms of $36, 24, 16, \dots$

5 Sum all terms of $0.4, 0.04, 0.004, 0.0004, \dots$

Infinite geometric series - exercises

Infinite geometric series

For $a_1, a_1r, a_1r^2, a_1r^3, \dots$:

$$\sum_{n=1}^{\infty} a_n = a_1 \cdot \frac{1}{1-r}$$

1 Find $\sum_{n=1}^{\infty} 6 \cdot \left(\frac{1}{3}\right)^{n-1}$

$$a_1 = 6, r = \frac{1}{3}$$

$$\Rightarrow \sum_{n=1}^{\infty} a_n = 6 \cdot \frac{1}{1-\frac{1}{3}} = \frac{6}{\frac{2}{3}} = 6 \cdot \frac{3}{2} = \frac{18}{2} = 9$$

2 Find $\sum_{n=1}^{\infty} 20 \cdot \left(\frac{2}{5}\right)^{n-1}$

$$a_1 = 20, r = \frac{2}{5}$$

$$\Rightarrow \sum_{n=1}^{\infty} a_n = 20 \cdot \frac{1}{1-\frac{2}{5}} = \frac{20}{\frac{3}{5}} = 20 \cdot \frac{5}{3} = \frac{100}{3}$$

3 Find $\sum_{n=1}^{\infty} 24 \cdot \left(-\frac{1}{2}\right)^{n-1}$

$$a_1 = 24, r = -\frac{1}{2}$$

Find the sum of the given geometric sequence.

1 Sum all terms of $12, 3, \frac{3}{4}, \frac{3}{16}, \dots$

2 Sum all terms of $-27, 9, -3, 1, \dots$

3 Sum all terms of $50, -30, 18, \dots$

4 Sum all terms of $36, 24, 16, \dots$

5 Sum all terms of $0.4, 0.04, 0.004, 0.0004, \dots$

Infinite geometric series - exercises

Infinite geometric series

For $a_1, a_1r, a_1r^2, a_1r^3, \dots$:

$$\sum_{n=1}^{\infty} a_n = a_1 \cdot \frac{1}{1-r}$$

1 Find $\sum_{n=1}^{\infty} 6 \cdot \left(\frac{1}{3}\right)^{n-1}$

$$a_1 = 6, r = \frac{1}{3}$$

$$\Rightarrow \sum_{n=1}^{\infty} a_n = 6 \cdot \frac{1}{1-\frac{1}{3}} = \frac{6}{\frac{2}{3}} = 6 \cdot \frac{3}{2} = \frac{18}{2} = 9$$

2 Find $\sum_{n=1}^{\infty} 20 \cdot \left(\frac{2}{5}\right)^{n-1}$

$$a_1 = 20, r = \frac{2}{5}$$

$$\Rightarrow \sum_{n=1}^{\infty} a_n = 20 \cdot \frac{1}{1-\frac{2}{5}} = \frac{20}{\frac{3}{5}} = 20 \cdot \frac{5}{3} = \frac{100}{3}$$

3 Find $\sum_{n=1}^{\infty} 24 \cdot \left(-\frac{1}{2}\right)^{n-1}$

$$a_1 = 24, r = -\frac{1}{2}$$

$$\Rightarrow \sum_{n=1}^{\infty} a_n = 24 \cdot \frac{1}{1-(-\frac{1}{2})} = \frac{24}{\frac{3}{2}} = 24 \cdot \frac{2}{3} = \frac{48}{3} = 16$$

Find the sum of the given geometric sequence.

1 Sum all terms of $12, 3, \frac{3}{4}, \frac{3}{16}, \dots$

2 Sum all terms of $-27, 9, -3, 1, \dots$

3 Sum all terms of $50, -30, 18, \dots$

4 Sum all terms of $36, 24, 16, \dots$

5 Sum all terms of $0.4, 0.04, 0.004, 0.0004, \dots$

Infinite geometric series - exercises

Infinite geometric series

For $a_1, a_1r, a_1r^2, a_1r^3, \dots$:

$$\sum_{n=1}^{\infty} a_n = a_1 \cdot \frac{1}{1-r}$$

1 Find $\sum_{n=1}^{\infty} 6 \cdot \left(\frac{1}{3}\right)^{n-1}$

$$a_1 = 6, r = \frac{1}{3}$$

$$\Rightarrow \sum_{n=1}^{\infty} a_n = 6 \cdot \frac{1}{1-\frac{1}{3}} = \frac{6}{\frac{2}{3}} = 6 \cdot \frac{3}{2} = \frac{18}{2} = 9$$

2 Find $\sum_{n=1}^{\infty} 20 \cdot \left(\frac{2}{5}\right)^{n-1}$

$$a_1 = 20, r = \frac{2}{5}$$

$$\Rightarrow \sum_{n=1}^{\infty} a_n = 20 \cdot \frac{1}{1-\frac{2}{5}} = \frac{20}{\frac{3}{5}} = 20 \cdot \frac{5}{3} = \frac{100}{3}$$

3 Find $\sum_{n=1}^{\infty} 24 \cdot \left(-\frac{1}{2}\right)^{n-1}$

$$a_1 = 24, r = -\frac{1}{2}$$

$$\Rightarrow \sum_{n=1}^{\infty} a_n = 24 \cdot \frac{1}{1-(-\frac{1}{2})} = \frac{24}{\frac{3}{2}} = 24 \cdot \frac{2}{3} = \frac{48}{3} = 16$$

Find the sum of the given geometric sequence.

1 Sum all terms of $12, 3, \frac{3}{4}, \frac{3}{16}, \dots$
 $a_1 = 12, r = \frac{1}{4}$

2 Sum all terms of $-27, 9, -3, 1, \dots$

3 Sum all terms of $50, -30, 18, \dots$

4 Sum all terms of $36, 24, 16, \dots$

5 Sum all terms of $0.4, 0.04, 0.004, 0.0004, \dots$

Infinite geometric series - exercises

Infinite geometric series

For $a_1, a_1r, a_1r^2, a_1r^3, \dots$:

$$\sum_{n=1}^{\infty} a_n = a_1 \cdot \frac{1}{1-r}$$

1 Find $\sum_{n=1}^{\infty} 6 \cdot \left(\frac{1}{3}\right)^{n-1}$

$$a_1 = 6, r = \frac{1}{3}$$

$$\Rightarrow \sum_{n=1}^{\infty} a_n = 6 \cdot \frac{1}{1-\frac{1}{3}} = \frac{6}{\frac{2}{3}} = 6 \cdot \frac{3}{2} = \frac{18}{2} = 9$$

2 Find $\sum_{n=1}^{\infty} 20 \cdot \left(\frac{2}{5}\right)^{n-1}$

$$a_1 = 20, r = \frac{2}{5}$$

$$\Rightarrow \sum_{n=1}^{\infty} a_n = 20 \cdot \frac{1}{1-\frac{2}{5}} = \frac{20}{\frac{3}{5}} = 20 \cdot \frac{5}{3} = \frac{100}{3}$$

3 Find $\sum_{n=1}^{\infty} 24 \cdot \left(-\frac{1}{2}\right)^{n-1}$

$$a_1 = 24, r = -\frac{1}{2}$$

$$\Rightarrow \sum_{n=1}^{\infty} a_n = 24 \cdot \frac{1}{1-(-\frac{1}{2})} = \frac{24}{\frac{3}{2}} = 24 \cdot \frac{2}{3} = \frac{48}{3} = 16$$

Find the sum of the given geometric sequence.

1 Sum all terms of $12, 3, \frac{3}{4}, \frac{3}{16}, \dots$

$$a_1 = 12, r = \frac{1}{4}$$

$$\Rightarrow \sum_{n=1}^{\infty} a_n = 12 \cdot \frac{1}{1-\frac{1}{4}} = \frac{12}{\frac{3}{4}} = 12 \cdot \frac{4}{3} = 16$$

2 Sum all terms of $-27, 9, -3, 1, \dots$

3 Sum all terms of $50, -30, 18, \dots$

4 Sum all terms of $36, 24, 16, \dots$

5 Sum all terms of $0.4, 0.04, 0.004, 0.0004, \dots$

Infinite geometric series - exercises

Infinite geometric series

For $a_1, a_1r, a_1r^2, a_1r^3, \dots$:

$$\sum_{n=1}^{\infty} a_n = a_1 \cdot \frac{1}{1-r}$$

- Find $\sum_{n=1}^{\infty} 6 \cdot \left(\frac{1}{3}\right)^{n-1}$
 $a_1 = 6, r = \frac{1}{3}$
 $\Rightarrow \sum_{n=1}^{\infty} a_n = 6 \cdot \frac{1}{1-\frac{1}{3}} = \frac{6}{\frac{2}{3}} = 6 \cdot \frac{3}{2} = \frac{18}{2} = 9$
- Find $\sum_{n=1}^{\infty} 20 \cdot \left(\frac{2}{5}\right)^{n-1}$
 $a_1 = 20, r = \frac{2}{5}$
 $\Rightarrow \sum_{n=1}^{\infty} a_n = 20 \cdot \frac{1}{1-\frac{2}{5}} = \frac{20}{\frac{3}{5}} = 20 \cdot \frac{5}{3} = \frac{100}{3}$
- Find $\sum_{n=1}^{\infty} 24 \cdot \left(-\frac{1}{2}\right)^{n-1}$
 $a_1 = 24, r = -\frac{1}{2}$
 $\Rightarrow \sum_{n=1}^{\infty} a_n = 24 \cdot \frac{1}{1-(-\frac{1}{2})} = \frac{24}{\frac{3}{2}} = 24 \cdot \frac{2}{3} = \frac{48}{3} = 16$

Find the sum of the given geometric sequence.

- Sum all terms of $12, 3, \frac{3}{4}, \frac{3}{16}, \dots$
 $a_1 = 12, r = \frac{1}{4}$
 $\Rightarrow \sum_{n=1}^{\infty} a_n = 12 \cdot \frac{1}{1-\frac{1}{4}} = \frac{12}{\frac{3}{4}} = 12 \cdot \frac{4}{3} = 16$
- Sum all terms of $-27, 9, -3, 1, \dots$
 $a_1 = -27, r = -\frac{1}{3}$
 $\Rightarrow \sum_{n=1}^{\infty} a_n = -27 \cdot \frac{1}{1-(-\frac{1}{3})} = \frac{-27}{\frac{4}{3}} = -27 \cdot \frac{3}{4} = -\frac{81}{4}$
- Sum all terms of $50, -30, 18, \dots$
- Sum all terms of $36, 24, 16, \dots$
- Sum all terms of $0.4, 0.04, 0.004, 0.0004, \dots$

Infinite geometric series - exercises

Infinite geometric series

For $a_1, a_1r, a_1r^2, a_1r^3, \dots$:

$$\sum_{n=1}^{\infty} a_n = a_1 \cdot \frac{1}{1-r}$$

- Find $\sum_{n=1}^{\infty} 6 \cdot \left(\frac{1}{3}\right)^{n-1}$
 $a_1 = 6, r = \frac{1}{3}$
 $\Rightarrow \sum_{n=1}^{\infty} a_n = 6 \cdot \frac{1}{1-\frac{1}{3}} = \frac{6}{\frac{2}{3}} = 6 \cdot \frac{3}{2} = \frac{18}{2} = 9$
- Find $\sum_{n=1}^{\infty} 20 \cdot \left(\frac{2}{5}\right)^{n-1}$
 $a_1 = 20, r = \frac{2}{5}$
 $\Rightarrow \sum_{n=1}^{\infty} a_n = 20 \cdot \frac{1}{1-\frac{2}{5}} = \frac{20}{\frac{3}{5}} = 20 \cdot \frac{5}{3} = \frac{100}{3}$
- Find $\sum_{n=1}^{\infty} 24 \cdot \left(-\frac{1}{2}\right)^{n-1}$
 $a_1 = 24, r = -\frac{1}{2}$
 $\Rightarrow \sum_{n=1}^{\infty} a_n = 24 \cdot \frac{1}{1-(-\frac{1}{2})} = \frac{24}{\frac{3}{2}} = 24 \cdot \frac{2}{3} = \frac{48}{3} = 16$

Find the sum of the given geometric sequence.

- Sum all terms of $12, 3, \frac{3}{4}, \frac{3}{16}, \dots$
 $a_1 = 12, r = \frac{1}{4}$
 $\Rightarrow \sum_{n=1}^{\infty} a_n = 12 \cdot \frac{1}{1-\frac{1}{4}} = \frac{12}{\frac{3}{4}} = 12 \cdot \frac{4}{3} = 16$
- Sum all terms of $-27, 9, -3, 1, \dots$
 $a_1 = -27, r = -\frac{1}{3}$
 $\Rightarrow \sum_{n=1}^{\infty} a_n = -27 \cdot \frac{1}{1-(-\frac{1}{3})} = \frac{-27}{\frac{4}{3}} = -27 \cdot \frac{3}{4} = -\frac{81}{4}$
- Sum all terms of $50, -30, 18, \dots$
 $a_1 = 50, r = \frac{-30}{50} = -\frac{3}{5}$
 $\Rightarrow \sum_{n=1}^{\infty} a_n = 50 \cdot \frac{1}{1-(-\frac{3}{5})} = \frac{50}{\frac{8}{5}} = 50 \cdot \frac{5}{8} = \frac{250}{8} = \frac{125}{4}$
- Sum all terms of $36, 24, 16, \dots$
- Sum all terms of $0.4, 0.04, 0.004, 0.0004, \dots$

Infinite geometric series - exercises

Infinite geometric series

For $a_1, a_1r, a_1r^2, a_1r^3, \dots$:

$$\sum_{n=1}^{\infty} a_n = a_1 \cdot \frac{1}{1-r}$$

- Find $\sum_{n=1}^{\infty} 6 \cdot \left(\frac{1}{3}\right)^{n-1}$
 $a_1 = 6, r = \frac{1}{3}$
 $\Rightarrow \sum_{n=1}^{\infty} a_n = 6 \cdot \frac{1}{1-\frac{1}{3}} = \frac{6}{\frac{2}{3}} = 6 \cdot \frac{3}{2} = \frac{18}{2} = 9$
- Find $\sum_{n=1}^{\infty} 20 \cdot \left(\frac{2}{5}\right)^{n-1}$
 $a_1 = 20, r = \frac{2}{5}$
 $\Rightarrow \sum_{n=1}^{\infty} a_n = 20 \cdot \frac{1}{1-\frac{2}{5}} = \frac{20}{\frac{3}{5}} = 20 \cdot \frac{5}{3} = \frac{100}{3}$
- Find $\sum_{n=1}^{\infty} 24 \cdot \left(-\frac{1}{2}\right)^{n-1}$
 $a_1 = 24, r = -\frac{1}{2}$
 $\Rightarrow \sum_{n=1}^{\infty} a_n = 24 \cdot \frac{1}{1-(-\frac{1}{2})} = \frac{24}{\frac{3}{2}} = 24 \cdot \frac{2}{3} = \frac{48}{3} = 16$

Find the sum of the given geometric sequence.

- Sum all terms of $12, 3, \frac{3}{4}, \frac{3}{16}, \dots$
 $a_1 = 12, r = \frac{1}{4}$
 $\Rightarrow \sum_{n=1}^{\infty} a_n = 12 \cdot \frac{1}{1-\frac{1}{4}} = \frac{12}{\frac{3}{4}} = 12 \cdot \frac{4}{3} = 16$
- Sum all terms of $-27, 9, -3, 1, \dots$
 $a_1 = -27, r = -\frac{1}{3}$
 $\Rightarrow \sum_{n=1}^{\infty} a_n = -27 \cdot \frac{1}{1-(-\frac{1}{3})} = \frac{-27}{\frac{4}{3}} = -27 \cdot \frac{3}{4} = -\frac{81}{4}$
- Sum all terms of $50, -30, 18, \dots$
 $a_1 = 50, r = \frac{-30}{50} = -\frac{3}{5}$
 $\Rightarrow \sum_{n=1}^{\infty} a_n = 50 \cdot \frac{1}{1-(-\frac{3}{5})} = \frac{50}{\frac{8}{5}} = 50 \cdot \frac{5}{8} = \frac{250}{8} = \frac{125}{4}$
- Sum all terms of $36, 24, 16, \dots$
 $a_1 = 36, r = \frac{24}{36} = \frac{2}{3}$
 $\Rightarrow \sum_{n=1}^{\infty} a_n = 36 \cdot \frac{1}{1-\frac{2}{3}} = \frac{36}{\frac{1}{3}} = 36 \cdot \frac{3}{1} = 108$
- Sum all terms of $0.4, 0.04, 0.004, 0.0004, \dots$

Infinite geometric series - exercises

Infinite geometric series

For $a_1, a_1r, a_1r^2, a_1r^3, \dots$:

$$\sum_{n=1}^{\infty} a_n = a_1 \cdot \frac{1}{1-r}$$

- Find $\sum_{n=1}^{\infty} 6 \cdot \left(\frac{1}{3}\right)^{n-1}$
 $a_1 = 6, r = \frac{1}{3}$
 $\Rightarrow \sum_{n=1}^{\infty} a_n = 6 \cdot \frac{1}{1-\frac{1}{3}} = \frac{6}{\frac{2}{3}} = 6 \cdot \frac{3}{2} = \frac{18}{2} = 9$
- Find $\sum_{n=1}^{\infty} 20 \cdot \left(\frac{2}{5}\right)^{n-1}$
 $a_1 = 20, r = \frac{2}{5}$
 $\Rightarrow \sum_{n=1}^{\infty} a_n = 20 \cdot \frac{1}{1-\frac{2}{5}} = \frac{20}{\frac{3}{5}} = 20 \cdot \frac{5}{3} = \frac{100}{3}$
- Find $\sum_{n=1}^{\infty} 24 \cdot \left(-\frac{1}{2}\right)^{n-1}$
 $a_1 = 24, r = -\frac{1}{2}$
 $\Rightarrow \sum_{n=1}^{\infty} a_n = 24 \cdot \frac{1}{1-(-\frac{1}{2})} = \frac{24}{\frac{3}{2}} = 24 \cdot \frac{2}{3} = \frac{48}{3} = 16$

Find the sum of the given geometric sequence.

- Sum all terms of $12, 3, \frac{3}{4}, \frac{3}{16}, \dots$
 $a_1 = 12, r = \frac{1}{4}$
 $\Rightarrow \sum_{n=1}^{\infty} a_n = 12 \cdot \frac{1}{1-\frac{1}{4}} = \frac{12}{\frac{3}{4}} = 12 \cdot \frac{4}{3} = 16$
- Sum all terms of $-27, 9, -3, 1, \dots$
 $a_1 = -27, r = -\frac{1}{3}$
 $\Rightarrow \sum_{n=1}^{\infty} a_n = -27 \cdot \frac{1}{1-(-\frac{1}{3})} = \frac{-27}{\frac{4}{3}} = -27 \cdot \frac{3}{4} = -\frac{81}{4}$
- Sum all terms of $50, -30, 18, \dots$
 $a_1 = 50, r = \frac{-30}{50} = -\frac{3}{5}$
 $\Rightarrow \sum_{n=1}^{\infty} a_n = 50 \cdot \frac{1}{1-(-\frac{3}{5})} = \frac{50}{\frac{8}{5}} = 50 \cdot \frac{5}{8} = \frac{250}{8} = \frac{125}{4}$
- Sum all terms of $36, 24, 16, \dots$
 $a_1 = 36, r = \frac{24}{36} = \frac{2}{3}$
 $\Rightarrow \sum_{n=1}^{\infty} a_n = 36 \cdot \frac{1}{1-\frac{2}{3}} = \frac{36}{\frac{1}{3}} = 36 \cdot \frac{3}{1} = 108$
- Sum all terms of $0.4, 0.04, 0.004, 0.0004, \dots$
 $a_1 = 0.4, r = 0.1$
 $\Rightarrow \sum_{n=1}^{\infty} a_n = 0.4 \cdot \frac{1}{1-0.1} = \frac{0.4}{0.9} = \frac{4}{9} = 0.444\bar{4}$

