

13.4 Nonlinear Systems

In this final section, we want to learn how to solve systems of equations that are not necessarily all linear. We call these non-linear systems of equations.

Definition: Non-linear system of equations

A system of equations where one or more equations involved is not a line.

We primarily use the substitution method to solve a non-linear system. However, sometimes elimination will work as well.

Also, just as before, the solution to a non-linear system is all the points of intersection of the graphs of the equations. Therefore, since we now have more than just lines, we can have a variety of numbers of solutions. The graph can intersect once, twice, several times or not at all. Therefore, we should always verify the solution(s) to a system by looking at the graph.

Example 1:

Solve the system.

a. $x^2 + y^2 = 100$
 $y - x = 2$

b. $x^2 + y^2 = 25$
 $y^2 = x + 5$

c. $x^2 + 2y^2 = 12$
 $xy = 4$

Solution:

a. We will solve this system by substitution. So we start by solving the bottom equation for y and then substitute it into the top equation. We get

$$y - x = 2 \Rightarrow y = x + 2$$

$$x^2 + y^2 = 100 \Rightarrow x^2 + (x + 2)^2 = 100$$

$$x^2 + x^2 + 4x + 4 = 100$$

$$2x^2 + 4x - 96 = 0$$

$$2(x^2 + 2x - 48) = 0$$

$$2(x + 8)(x - 6) = 0$$

$$x = -8, 6$$


So we have two different x values. This means we should have two points of intersection. Let's now find the y values and verify with a graph.

$$y = x + 2 \qquad y = x + 2$$

$$y = -8 + 2 \qquad y = 6 + 2$$

$$y = -6 \qquad y = 8$$

So our solutions are $(-8, -6)$ and $(6, 8)$. We clearly have a circle and a line, thus we can easily graph them together and get



- b. **Again we will use substitution to solve.** This time notice that the bottom equation is already solved for y^2 and we have a y^2 in the top equation. Thus, that is the substitution we will make. We get

$$\begin{aligned} x^2 + y^2 &= 25 \\ \underbrace{}_{y^2 = x + 5} &\Rightarrow x^2 + (x + 5) = 25 \\ y^2 &= x + 5 \end{aligned}$$


Now we solve for x and then solve for y.

$$\begin{aligned} x^2 + (x + 5) &= 25 \\ x^2 + x - 20 &= 0 \\ (x + 5)(x - 4) &= 0 \\ x &= -5, 4 \end{aligned}$$

We substitute these back in to get y. We get

$$\begin{aligned} y^2 &= -5 + 5 & y^2 &= 4 + 5 \\ y^2 &= 0 & y^2 &= 9 \\ y &= 0 & y &= \pm 3 \end{aligned}$$

So we have three different solutions $(-5, 0)$, $(4, 3)$ and $(4, -3)$. Lets verify with a graph. We have here a parabola and a circle. We get



- c. **Again, we will use substitution to solve.** We need to decide which variable to solve for first. It seems that x or y on the bottom equation would be easiest. So we will solve for x. We get

$$xy = 4 \Rightarrow x = \frac{4}{y}$$

Now we substitute that into the first equation and solve. We have

$$\begin{aligned} x^2 + 2y^2 = 12 &\Rightarrow \left(\frac{4}{y}\right)^2 + 2y^2 = 12 \\ \frac{16}{y^2} + 2y^2 &= 12 \end{aligned}$$

We will have to clear the fractions and solve as we did in chapter 10, that is, using a substitution.

$$y^2 \left(\frac{16}{y^2} + 2y^2 \right) = y^2(12)$$

$$16 + 2y^4 = 12y^2$$

$$2y^4 - 12y^2 + 16 = 0$$

Substitute $u = y^2$

$$2u^2 - 12u + 16 = 0$$

$$2(u^2 - 6u + 8) = 0$$

$$2(u - 4)(u - 2) = 0$$

$$u = 4, 2$$

Re-substitute $u = y^2$

$$y^2 = 4 \quad y^2 = 2$$

$$y = \pm 2 \quad y = \pm\sqrt{2}$$

So since we have four different y values we will have to find x for each one. We

substitute these in to $x = \frac{4}{y}$ to get

$$x = \frac{4}{2}$$

$$= 2$$

$$x = \frac{4}{-2}$$

$$= -2$$

$$x = \frac{4}{\sqrt{2}} \cdot \frac{\sqrt{2}}{\sqrt{2}}$$

$$= \frac{4\sqrt{2}}{2}$$

$$= 2\sqrt{2}$$

$$x = \frac{4}{-\sqrt{2}} \cdot \frac{\sqrt{2}}{\sqrt{2}}$$


$$= -\frac{4\sqrt{2}}{2}$$

$$= -2\sqrt{2}$$

So we have four solutions, $(2, 2)$, $(-2, -2)$, $(2\sqrt{2}, \sqrt{2})$ and $(-2\sqrt{2}, -\sqrt{2})$.

Lets verify with a graph. We have an ellipse and a basic function ($xy = 4 \Rightarrow y = \frac{4}{x}$).

So we have



Example 2:

Solve the system.

a. $y = 3^x$
 $y = 3^{2x} - 2$

b. $y = \log_2(x+1)$
 $y = 5 - \log_2(x-3)$

Solution:

- a. This time solving is a little more complicated. There are a variety of direction we could go, however, we are going to start by noticing $3^{2x} = (3^x)^2$. So our system really is

$$y = 3^x$$

$$y = (3^x)^2 - 2$$

We can see clearly we will substitute the top equation into the bottom. That is, put y in for 3^x . We get

$$y = (y)^2 - 2$$

$$y = y^2 - 2$$

$$y^2 - y - 2 = 0$$

$$(y - 2)(y + 1) = 0$$

$$y = 2, -1$$

Now we substitute these values back in to get

$$2 = 3^x$$

$$-1 = 3^x$$

$$x = \log_3 2$$

However, the second equation is impossible (recall, exponential functions are always positive). Thus, that solution must be omitted. So we have a solution of $(\log_2 3, 2)$.

- b. Lastly, since these equations are both already solved for y , we can simply set them equal to one another. Then we are left with a logarithmic equation to solve. We get

$$\log_2(x+1) = 5 - \log_2(x-3)$$

$$\log_2(x+1) + \log_2(x-3) = 5$$

$$\log_2(x+1)(x-3) = 5$$

$$(x+1)(x-3) = 2^5$$

$$x^2 - 2x - 3 = 32$$

$$x^2 - 2x - 35 = 0$$

$$(x-7)(x+5) = 0$$

$$x = 7, -5$$

However, -5 cannot be a solution since it doesn't even check in the equation. Thus we only have $x = 7$. Now we substitute this back into either original equation to get the y value. We choose the first equation.

$$y = \log_2(7+1)$$

$$= \log_2 8$$

$$= 3$$

So the solution is $(7, 3)$.

13.4 Exercises

Solve the systems.

1. $x^2 + y^2 = 2$
 $x + y = 2$

2. $x^2 + y^2 = 25$
 $y - x = 1$

3. $25x^2 + 9y^2 = 225$
 $5x + 3y = 15$

4. $9x^2 + 4y^2 = 36$
 $3x + 2y = 6$

5. $y^2 = x + 3$
 $2y = x + 4$

6. $y = x^2$
 $3x = y + 2$

7. $x^2 - y^2 = 16$
 $x - 2y = 1$

8. $x^2 + 4y^2 = 25$
 $x + 2y = 7$

9. $x^2 + y^2 = 18$
 $2x + y = 3$

10. $x^2 - y = 3$
 $2x - y = 3$

11. $x^2 + y^2 = 20$
 $y = x^2$

12. $x^2 - y^2 = 3$
 $y = x^2 - 3$

13. $x^2 - x - y = 2$
 $4x - 3y = 0$

14. $x^2 - 2x + 2y^2 = 8$
 $2x + y = 6$

15. $x^2 + y^2 = 13$
 $y = x^2 - 1$

16. $x^2 - y = 5$
 $x^2 + y^2 = 25$

17. $x^2 + y^2 = 25$
 $2x^2 - 3y^2 = 5$

18. $x^2 + y^2 = 4$
 $9x^2 + 16y^2 = 144$

19. $x^2 + y^2 = 13$
 $x^2 - y^2 = -16$

20. $x^2 + y^2 = 16$
 $y^2 - 2y^2 = 10$

21. $x^2 + y^2 = 20$
 $x^2 - y^2 = -12$

22. $x^2 + y^2 = 14$
 $x^2 - y^2 = 4$

23. $xy = -\frac{9}{5}$
 $3x + 2y = 6$

24. $x + y = -6$
 $xy = -7$

25. $y = x^2 - 4$
 $x^2 - y^2 = -16$

26. $x^2 + y^2 = 25$
 $y^2 = x + 5$

27. $xy = \frac{1}{6}$
 $y + x = 5xy$

28. $xy = \frac{1}{12}$
 $x + y = 7xy$

29. $x^2 + xy + 2y^2 = 7$
 $x - 2y = 5$

30. $x^2 - xy + 3y^2 = 27$
 $x - y = 2$

31. $x^2 + y^2 = 5$
 $xy = 2$

32. $x^2 + y^2 = 20$
 $xy = 8$

33. $3xy + x^2 = 34$
 $2xy - 3x^2 = 8$

34. $2xy + 3y^2 = 7$
 $3xy - 2y^2 = 4$

35. $\frac{1}{x} + \frac{1}{y} = 5$
 $\frac{1}{x} - \frac{1}{y} = -3$

36. $\frac{1}{x} - \frac{1}{y} = 4$
 $\frac{1}{x} + \frac{1}{y} = -2$

$$37. \begin{aligned} \frac{2}{x^2} + \frac{5}{y^2} &= 3 \\ \frac{3}{x^2} - \frac{2}{y^2} &= 1 \end{aligned}$$

$$38. \begin{aligned} \frac{1}{x^2} - \frac{3}{y^2} &= 14 \\ \frac{2}{x^2} + \frac{1}{y^2} &= 35 \end{aligned}$$

$$39. \begin{aligned} x^4 &= y - 1 \\ y - 3x^2 + 1 &= 0 \end{aligned}$$

$$40. \begin{aligned} x^3 - y &= 0 \\ xy - 16 &= 0 \end{aligned}$$

$$41. \begin{aligned} y &= -\sqrt{x} \\ (x-3)^2 + y^2 &= 4 \end{aligned}$$

$$42. \begin{aligned} y &= \sqrt{x} \\ (x+2)^2 + y^2 &= 1 \end{aligned}$$

$$43. \begin{aligned} y &= 2^x \\ y &= 2^{2x} - 12 \end{aligned}$$

$$44. \begin{aligned} y &= 5^x \\ y &= 5^{2x} - 1 \end{aligned}$$

$$45. \begin{aligned} y &= e^{4x} \\ y &= e^{2x} + 6 \end{aligned}$$

$$46. \begin{aligned} y &= 2e^{2x} \\ y &= e^x - 1 \end{aligned}$$

$$47. \begin{aligned} y &= \log_2(x+4) \\ y &= 2 - \log_2(x+1) \end{aligned}$$

$$48. \begin{aligned} y &= \log_6 x \\ y &= -\log_6(x+1) \end{aligned}$$

$$49. \begin{aligned} y &= \log_9(x+1) \\ y &= \log_9 x + \frac{1}{2} \end{aligned}$$

$$50. \begin{aligned} y &= \log_{16}(x+3) \\ y &= \log_{16}(x-1) + \frac{1}{2} \end{aligned}$$