



① $y = 2 \sin(3x - \pi)$ $A = 2$ $B = 3$ $C = -\pi$
 Amplitude = 2 phase shift = $-\frac{C}{B} = \frac{\pi}{3}$
 Period = $|\frac{2\pi}{B}| = \frac{2\pi}{3}$

cycle starts at $\frac{\pi}{3}$ (phase shift) and ends at $\frac{\pi}{3} + \frac{2\pi}{3} = \pi$ (phase shift + period)



② $y = -5 \cos(x + \frac{\pi}{2})$ $A = -5$ $B = 1$ $C = \frac{\pi}{2}$
 Amplitude = $|A| = 5$ phase shift = $-\frac{C}{B} = -\frac{\pi}{2}$
 Period = $\frac{2\pi}{1} = 2\pi$

cycle starts at $-\frac{\pi}{2}$ and ends at $-\frac{\pi}{2} + 2\pi = \frac{3\pi}{2}$



③ $-2 \log x + \frac{1}{2} \log y - 4 \log z = -\log x^2 + \log y^{\frac{1}{2}} - \log z^4 =$
 $-\log x^2 + \log \sqrt{y} - \log z^4 = \boxed{\log \left(\frac{\sqrt{y}}{x^2 z^4} \right)}$

④ $P(t) = 35.7(1+0.01)^t$ where $t=0$ is the year 2010
 population in 2015 = $P(5) = 35.7(1.01)^5 = 37.52105879 = \boxed{37.5 \text{ million}}$
 $35.7 \times 2 = 71.4$. We need to find in what year the population will be 71.4 million

$71.4 = 35.7(1.01)^t$ divide by 35.7

$2 = (1.01)^t$ Now apply log or ln

$\ln 2 = \ln(1.01)^t$

$\ln 2 = t \ln(1.01)$ $t = \frac{\ln 2}{\ln(1.01)} = 69.66071689$

So the population will be double sometimes in the **year 2079**

⑤ $f(t) = 200(1-.12)^t$ where $t=0$ is the year 2012

So $f(t) = 200(.88)^t$

a) $t=7$ $f(7) = 200(.88)^7 = 81.73511972 = \boxed{81.7 \text{ grams in 2019}}$

b) $50 = 200(.88)^t$ divide by 200

$\frac{50}{200} = (.88)^t$ $\frac{1}{4} = (.88)^t$ Now apply log or ln

$\ln(\frac{1}{4}) = \ln(.88)^t$

$\ln(\frac{1}{4}) = t \ln(.88)$ $t = \frac{\ln(\frac{1}{4})}{\ln(.88)} = 10.84454196$

So there will be 50 grams left **sometimes in the year 2022**

⑥ $\log_3(x) + \log_3(x-8) = 2$

$\log_3(x(x-8)) = 2$

$\log_3(x^2 - 8x) = 2$

$x^2 - 8x = 3^2 = 9$

$x^2 - 8x = 9$

$x^2 - 8x - 9 = 0$

$(x-9)(x+1) = 0$

$x=9$

~~$x=-1$~~ reject since inputs of log are positive

⑦ $f(x) = -\log(3-2x)$

Domain $3-2x > 0$

$-2x > -3$

$x < \frac{3}{2}$


$(-\infty, \frac{3}{2})$

v. A. $x = \frac{3}{2}$

x-intercept $3-2x = 1$

$-2x = -2$

$x=1$ $(1,0)$



$$\textcircled{8} \quad \ln \sqrt{\frac{xy^3}{z}} = \ln \left(\frac{xy^3}{z^{1/2}} \right)^{1/2} = \ln \left(\frac{x^{1/2} y^{3/2}}{z^{1/4}} \right)$$


$$= \ln x^{1/2} + \ln y^{3/2} - \ln z^{1/4} = \frac{1}{2} \ln x + \frac{3}{2} \ln y - \frac{1}{4} \ln z = \boxed{\frac{1}{2} u + \frac{3}{2} v - \frac{1}{4} w}$$

$$\textcircled{9} \quad \tan(\alpha) = -\frac{5}{12} = \frac{O}{A}$$

$$5^2 + (-12)^2 = R^2$$

$$25 + 144 = 169 \quad R=13$$

$$\sin(\alpha) = \frac{5}{13} \quad \cos(\alpha) = -\frac{12}{13}$$



$$\sin(2\alpha) = 2 \sin \alpha \cos \alpha = 2 \cdot \frac{5}{13} \cdot \left(-\frac{12}{13}\right) = \boxed{-\frac{120}{169}}$$

$$\cos(2\alpha) = \cos^2 \alpha - \sin^2 \alpha = \left(-\frac{12}{13}\right)^2 - \left(\frac{5}{13}\right)^2 = \frac{144 - 25}{169} = \boxed{\frac{119}{169}}$$

$$\textcircled{10} \quad \text{a) } \tan^{-1}\left(\frac{\sqrt{3}}{3}\right) = \frac{\pi}{6}$$

$$\boxed{x = \frac{\pi}{6} + N\pi \quad N=0, \pm 1, \pm 2, \dots}$$

$$\text{b) } \cos^{-1}(-1) = \pi$$

$$\boxed{x = \pi + 2N\pi \quad N=0, \pm 1, \pm 2, \dots}$$

$$\text{c) } \sin^{-1}\left(-\frac{\sqrt{3}}{2}\right) = -\frac{\pi}{3} \quad \text{solution 1}$$

$$\pi - \left(-\frac{\pi}{3}\right) = \pi + \frac{\pi}{3} = \frac{4\pi}{3} \quad \text{solution 2}$$

general solution

$$\boxed{x = -\frac{\pi}{3} + 2N\pi \quad N=0, \pm 1, \pm 2, \dots}$$

$$\boxed{x = \frac{4\pi}{3} + 2N\pi \quad N=0, \pm 1, \pm 2, \dots}$$

$$\text{d) } 2 \cos^2 x = \cos x$$

$$2 \cos^2 x - \cos x = 0$$

$$\cos x (2 \cos x - 1) = 0$$

$$\cos x = 0$$

$$\text{gives } x = \pm \cos^{-1}(0) + 2N\pi$$

$$x = \pm \frac{\pi}{2} + 2N\pi$$

$$N = 0, \pm 1, \pm 2$$

$$2 \cos x = 1 \quad \cos x = \frac{1}{2}$$

$$x = \pm \cos^{-1}\left(\frac{1}{2}\right) + 2N\pi$$

$$x = \pm \frac{\pi}{3} + 2N\pi$$

$$N = 0, \pm 1, \pm 2$$

general solution

$$\boxed{x = \pm \frac{\pi}{2} + 2N\pi \quad N=0, \pm 1, \pm 2, \dots}$$

$$\boxed{x = \pm \frac{\pi}{3} + 2N\pi \quad N=0, \pm 1, \pm 2, \dots}$$

$$\text{e) } 2 \sin^2 x + \sin x - 1 = 0. \quad \text{Let } \sin x = u$$

$$2u^2 + u - 1 = 0 \quad \text{gives}$$

$$u = \frac{-1 \pm \sqrt{1 - 4(2)(-1)}}{4} = \frac{-1 \pm 3}{4} \quad \begin{matrix} -1 \\ \frac{1}{2} \end{matrix}$$

$$\sin x = -1$$

$$\sin x = \frac{1}{2}$$

$$\boxed{x = -\frac{\pi}{2} + 2N\pi \quad N=0, \pm 1, \pm 2, \dots}$$

$$\boxed{x = \frac{\pi}{6} + 2N\pi \quad N=0, \pm 1, \pm 2, \dots}$$

$$\boxed{x = \frac{5\pi}{6} + 2N\pi \quad N=0, \pm 1, \pm 2, \dots}$$

$$(11) \quad 3e^{2x+7} = 63 \quad \text{divide by 3}$$

$$e^{2x+7} = 21$$

$$\ln(e^{2x+7}) = \ln(21)$$

$$(2x+7) \ln e = \ln(21)$$

Recall $\ln e = 1$

$$2x+7 = \ln(21)$$

$$2x = \ln(21) - 7$$

$$x = \frac{\ln(21) - 7}{2}$$

exact answer

$$x = -1.977738781$$

$$x = -1.98$$

approximation