

MAT 1575 Final Exam Review Problems

Revised by Prof. Kostadinov Spring 2014, Prof. ElHitti Summer 2017, Prof. Africk Fall 2019

1. Evaluate the following definite integrals:

$$\text{a. } \int_0^1 x^2(x^3 + 1)^3 dx \quad \text{b. } \int_0^1 \frac{x}{\sqrt{x^2 + 9}} dx \quad \text{c. } \int_0^1 \frac{3x^2}{\sqrt[3]{x^3 + 1}} dx$$

2. Evaluate the following indefinite integrals:

$$\text{a. } \int x^2 \ln(x) dx \quad \text{b. } \int x^2 e^{-x} dx \quad \text{c. } \int x \cos(3x) dx$$

3. Find the area of the region enclosed by the graphs of:

$$\text{a. } y = 3 - x^2 \text{ and } y = -2x \quad \text{b. } y = x^2 - 2x \text{ and } y = x + 4$$

4. Find the volume of the solid obtained by rotating the region bounded by the graphs of:

$$\text{a. } y = x^2 - 9, y = 0 \text{ about the x-axis.} \quad \text{b. } y = 16 - x, y = 3x + 12, x = -1 \text{ about the x-axis.}$$

$$\text{c. } y = x^2 + 2, y = -x^2 + 10, x \geq 0 \text{ about the y-axis.}$$

5. Evaluate the following indefinite integrals:

$$\text{a. } \int \frac{1}{x^2 \sqrt{36 - x^2}} dx \quad \text{b. } \int \frac{\sqrt{x^2 - 9}}{x^4} dx \quad \text{c. } \int \frac{9}{x^2 \sqrt{x^2 + 9}} dx \quad \text{d. } \int \frac{6}{x^2 \sqrt{x^2 - 36}} dx$$

6. Evaluate the following indefinite integrals:

$$\text{a. } \int \frac{3x+7}{x^2+6x+9} dx \quad \text{b. } \int \frac{5x+6}{x^2-36} dx \quad \text{c. } \int \frac{3x+2}{x^2+2x-8} dx \quad \text{d. } \int \frac{12-8x}{x^2(x-6)} dx \quad \text{e. } \int \frac{-2x^2+4x+4}{x(x-2)^2} dx$$

7. Evaluate the improper integral:

$$\text{a. } \int_0^{\infty} \frac{2}{(x+2)^3} dx \quad \text{b. } \int_0^{\infty} \frac{5}{\sqrt[5]{x+5}} dx \quad \text{c. } \int_3^5 \frac{3}{\sqrt[3]{(x-3)^4}} dx$$

8. Decide if the following series converges or not. Justify your answer using an appropriate test:

$$\text{a. } \sum_{n=1}^{\infty} \frac{9n^5}{3n^5+5} \quad \text{b. } \sum_{n=1}^{\infty} \frac{5}{10^n} \quad \text{c. } \sum_{n=1}^{\infty} \frac{5n}{10^n} \quad \text{d. } \sum_{n=1}^{\infty} \frac{n!}{n^2 5^n} \quad \text{e. } \sum_{n=1}^{\infty} \left(\frac{n+1}{2n+3} \right)^n$$

9. Determine whether the series is absolutely or conditionally convergent or divergent:

$$\text{a. } \sum_{n=1}^{\infty} (-1)^n \frac{10}{7n+2} \quad \text{b. } \sum_{n=1}^{\infty} \frac{(-1)^n}{\sqrt{n^5}} \quad \text{c. } \sum_{n=0}^{\infty} (-1)^n 5^{-n} \quad \text{d. } \sum_{n=1}^{\infty} (-1)^n \frac{n^2 - n - 1}{2n^2 + n + 1}$$

10. Find the radius and the interval of convergence of the following power series:

$$a. \sum_{n=0}^{\infty} \frac{(x-1)^n}{n+2} \quad b. \sum_{n=0}^{\infty} \frac{(-1)^n(x-1)^n}{n+2} \quad c. \sum_{n=1}^{\infty} \frac{(x+1)^n}{n5^n} \quad d. \sum_{n=1}^{\infty} \frac{(-1)^n(x+1)^n}{n5^n}$$

11. Find the Taylor polynomial of degree 2 for the given function, centered at the given number a :

a. $f(x) = e^{-2x}$ at $a = -1$. b. $f(x) = \cos(5x)$ at $a = 2\pi$.


12. Find the Taylor polynomial of degree 3 for the given function, centered at the given number a :

a. $f(x) = 1 + e^{-x}$ at $a = -1$ b. $f(x) = \sin(x)$ at $a = \frac{\pi}{2}$

Answers:


(1a). $\frac{5}{4}$ (1b). $\sqrt{10} - 3$ (1c). $\frac{3}{2} (2^{\frac{2}{3}} - 1)$

(2a). $\frac{x^3 \ln(x)}{3} - \frac{x^3}{9} + C$ (2b). $-(x^2 + 2x + 2)e^{-x} + C$ (2c). $\frac{1}{3}x \sin(3x) + \frac{1}{9} \cos(3x) + C$




(3a). The area of the region between the two curves is:

$$Area = \int_{-1}^3 (3 - x^2 - (-2x)) dx = \frac{32}{3}$$



(3b). The area of the region between the two curves is:

$$Area = \int_{-1}^5 (x + 4 - (x^2 - 2x)) dx = \frac{125}{6}$$



(4a). Approximate the volume of the solid by vertical disks with radius $y = x^2 - 9$ between $x = -3$ and $x = 3$; gives the volume is $V = \int_{-3}^3 \pi(x^2 - 9)^2 dx = \frac{1296}{5} \pi$.

(4b). Using a washer of outer radius $R_{outer} = 16 - x$ and inner radius $R_{inner} = 3x + 12$ at x , gives the volume:

$$V = \pi \int_{-1}^1 ((16 - x)^2 - (3x + 12)^2) dx = \frac{656\pi}{3}$$

where the upper limit 1 is obtained from $16 - x = 3x + 12 \Rightarrow x = 1$.

(4c). 16π

(5a). $-\frac{\sqrt{36-x^2}}{36x} + C$ (5b). $\frac{(x^2-9)^{3/2}}{27x^3} + C$ (5c). $-\frac{\sqrt{x^2+9}}{x} + C$ (5d). $\frac{\sqrt{x^2-36}}{6x} + C$

(6a). $\frac{2}{x+3} + 3\ln|x+3| + C$ (6b). $3\ln|x-6| + 2\ln|x+6| + C$ (6c). $\frac{5}{3}\ln|x+4| + \frac{4}{3}\ln|x-2| + C$

(6d). $\frac{2}{x} - \ln|x-6| + \ln|x| + C$ (6e). $\ln|x| - \frac{2}{x-2} - 3\ln|x-2| + C$

(7a). $\frac{1}{4}$ (7b). The integral does not converge (7c). The integral does not converge

(8a). $\lim_{n \rightarrow \infty} \frac{9n^5}{3n^5+5} = \lim_{n \rightarrow \infty} \frac{3}{1+5/3n^5} = 3 > 0$ so the series diverges by the nth term test for divergence.

(8b). This is a geometric series, with common ratio $r = 1/10 < 1$, so it converges to $5/9$:

$$\sum_{n=1}^{\infty} \frac{5}{10^n} = \frac{a}{1-r} = \frac{5/10}{1-\frac{1}{10}} = \frac{5/10}{9/10} = \frac{5}{9}$$

(8c). $\lim_{n \rightarrow \infty} \frac{5(n+1)}{10^{n+1}} / \frac{5n}{10^n} = \lim_{n \rightarrow \infty} \frac{1+1/5n}{10} = \frac{1}{10} < 1$ so the series converges by the ratio test.

(8d). $\lim_{n \rightarrow \infty} \frac{(n+1)!}{(n+1)^2 5^{n+1}} / \frac{n!}{n^2 5^n} = \lim_{n \rightarrow \infty} \frac{n^2}{5(n+1)} = \lim_{n \rightarrow \infty} \frac{n}{5(1+\frac{1}{n})} = \lim_{n \rightarrow \infty} \frac{n}{5} = \infty$ so the series diverges by the ratio test.

(8e). $\lim_{n \rightarrow \infty} \left[\left(\frac{n+1}{2n+3} \right)^n \right]^{\frac{1}{n}} = \lim_{n \rightarrow \infty} \frac{n+1}{2n+3} = \lim_{n \rightarrow \infty} \frac{1+\frac{1}{n}}{2+\frac{3}{n}} = \frac{1}{2} < 1$ so the series converges by the nth root test.

(9a). Conditionally convergent: The series converges by the alternating series test since

$\frac{10}{7n+2} > \frac{10}{7(n+1)+2}$ and $\lim_{n \rightarrow \infty} \frac{10}{7n+2} = 0$ but not absolutely since $\sum_{n=1}^{\infty} \left| (-1)^n \frac{10}{7n+2} \right| = \sum_{n=1}^{\infty} \frac{10}{7n+2}$ diverges by comparing it with $\sum_{n=1}^{\infty} \left| (-1)^n \frac{1}{\sqrt{n^5}} \right| = \sum_{n=1}^{\infty} \frac{1}{n^{5/2}}$ which diverges, using the limit comparison test: $\lim_{n \rightarrow \infty} \frac{10}{7n+2} / \frac{1}{n} = \lim_{n \rightarrow \infty} \frac{10n}{7n+2} = \frac{10}{7} > 0$

(9b). Absolutely convergent:

a convergent p-series with $p = 5/2 > 1$.

(9c). Absolutely convergent: $\sum_{n=0}^{\infty} |(-1)^n 5^{-n}| = \sum_{n=0}^{\infty} 5^{-n}$ is a convergent geometric series with common ratio $r = 1/5 < 1$.

(9d). $\lim_{n \rightarrow \infty} \frac{n^2 - n - 1}{2n^2 + n + 1} = \lim_{n \rightarrow \infty} \frac{1 - \frac{1}{n} - \frac{1}{n^2}}{2 + \frac{1}{n} + \frac{1}{n^2}} = \frac{1}{2} > 0$ so the series diverges by the nth term test for divergence.

(10a). The power series converges when $|x - 1| < 1$ by the ratio test, which gives a radius of convergence 1 and interval of convergence centered at 1. The series diverges at $x = 2$ (harmonic series) but converges at $x = 0$ (alternate harmonic series), so the interval of convergence is $0 \leq x < 2$.

(10b). The power series converges when $|x - 1| < 1$ by the ratio test, which gives a radius of convergence 1 and interval of convergence centered at 1. The series diverges at $x = 0$ (harmonic series) but converges at $x = 2$ (alternate harmonic series), so the interval of convergence is $0 < x \leq 2$.

(10c). The power series converges when $|x + 1| < 5$ by the ratio test, which gives a radius of convergence 5 and interval of convergence centered at -1 . The series diverges at $x = 4$ (harmonic series) but converges at $x = -6$ (alternate harmonic series), so the interval of convergence is $-6 \leq x < 4$.

(10d). The power series converges when $|x + 1| < 5$ by the ratio test, which gives a radius of convergence 5 and interval of convergence centered at -1 . The series diverges at $x = -6$ (harmonic series) but converges at $x = 4$ (alternate harmonic series), so the interval of convergence is $-6 < x \leq 4$.

$$(11a). \quad p_2(x) = e^2 - 2e^2(x + 1) + 2e^2(x + 1)^2$$

$$(11b). \quad p_2(x) = 1 - \frac{25}{2} (x - 2\pi)^2$$

$$(12a). \quad p_3(x) = 1 + e - e(x + 1) + \frac{e}{2}(x + 1)^2 - \frac{e}{6}(x + 1)^3$$

$$= 1 + \frac{e}{3} - \frac{e}{2}x - \frac{e}{6}x^3$$

$$(12b). \quad p_3(x) = 1 - \frac{1}{2} (x - \frac{\pi}{2})^2$$