

Question:	1	2	3	4	5	Total
Points:	5	5	5	5	5	25
Score:						

1. (5 points) Suppose an oil tanker starts leaking oil, creating an expanding circular oil spill on the water.

(a) Draw a picture illustrating the situation, and label the radius of the oil spill with the variable $r(t)$:

Solution: Just draw a circle, with maybe an oil tanker at the center, and label the radius $r(t)$. This video of a similar related rates exercise includes a picture:
<https://www.youtube.com/watch?app=desktop&v=nh56AppTg6U>

(b) Express the surface area $A(t)$ of the oil spill as a function of its radius $r(t)$:

Solution: Area of a circle as a function of its radius: $A(t) = \pi r(t)^2$

(c) Clearly, if oil spill is expanding, both the surface area and the radius are increasing with time. Use the Chain Rule to express *the rate of change* of the surface area $A(t)$ in terms of the radius $r(t)$ and the rate of change of the radius:

Solution:

$$\frac{dA}{dt} = 2\pi \cdot r(t) \cdot \frac{dr}{dt}$$

(d) Now answer the questions in Problem 1 of the WebWork set “Exam 3,” showing all your calculations below. Include units in your calculations:

Solution: WebWork randomizes the numbers in a given exercise. Here is an example:
 “The radius of a circular oil slick expands at a rate of 4 m/min.”
 The rate at which the radius is increasing is the value of $\frac{dr}{dt}$. So in this case: $\frac{dr}{dt} = 4m/min$.
 “How fast is the area of the oil slick increasing when the radius is 27 m?”
 We need to calculate $\frac{dA}{dt}$ when $r = 27$; we substitute into the “differential equation” we found in (b):

$$\left. \frac{dA}{dt} \right|_{r=27m} = 2\pi \cdot 27(m) \cdot 4(m/min) = 216\pi m^2/min$$

“If the radius is 0 at time $t = 0$, how fast is the area increasing after 2 mins?” We need to find the radius after 2 mins. Since $r(0) = 0$, and since the radius increases at 4 m/min, $r(2) = 0m + 2(mins) \cdot 4(m/min) = 8m$, i.e., the radius is 8m after 2 mins. We substitute this value of r into the equation:

$$\left. \frac{dA}{dt} \right|_{r=8m} = 2\pi \cdot 8(m) \cdot 4(m/min) = 64\pi m^2/min$$

Entered	Answer Preview	Result
678.584013175395	216π	correct
201.061929829747	64π	correct

All of the answers above are correct.

The radius of a circular oil slick expands at a rate of 4 m/min.

(a) How fast is the area of the oil slick increasing when the radius is 27 m?

$\frac{dA}{dt} = 216 \cdot \pi m^2/min$

(b) If the radius is 0 at time $t = 0$, how fast is the area increasing after 2 mins?

$\frac{dA}{dt} = 64\pi m^2/min$

2. (5 points) Recall that the “linearization” (or “(local) linear approximation”) $L(x)$ for a given function $f(x)$ at a point $x = x_0$ is just the equation of the tangent line for the graph $y = f(x)$ at the point $(x_0, f(x_0))$.

(a) Look at Problem 2 in the WebWork set “Exam 3.” As asked in part (a) of the WebWork exercise, find the linear approximation $L(x)$ for the given function $f(x)$ at the given value of x_0 , via writing down the following:

Solution: Again, every WebWork exercise will have different numbers. Suppose $f(x) = \sqrt{5+x}$ and $x_0 = 4$. Then:

$$f(x_0) = f(4) = \sqrt{9} = 3$$

$$f'(x) = \frac{1}{2\sqrt{x+9}}$$

$$f'(0) = \frac{1}{2\sqrt{9}} = \frac{1}{6}$$

$$L(x) = 3 + \frac{1}{6}(x - 4)$$

(b) Use the linear approximation from part (a) to estimate the values of $f(x)$ asked for in parts (b) and (c) of the WebWork exercise. Show your calculations here (you should not need a calculator for this!):


Solution: Suppose we are asked to use the linear approximation to approximate $\sqrt{8.9}$ and $\sqrt{9.1}$. In order to use our linear approximation to $f(x) = \sqrt{5+x}$, we have to express these values in terms of $f(x)$: note that $\sqrt{9.1} = \sqrt{5+4.1} = f(4.1)$. So we calculate $L(4.1)$:

$$\sqrt{9.1} = \sqrt{5+4.1} = f(4.1) \approx L(4.1) = 3 + \frac{1}{6}(4.1 - 4) = 3 + \frac{1}{6}(0.1) = 3 + \frac{1}{60}$$

Similarly:

$$\sqrt{8.9} = \sqrt{5+3.9} = f(3.9) \approx L(3.9) = 3 + \frac{1}{6}(3.9 - 4) = 3 + \frac{1}{6}(-0.1) = 3 - \frac{1}{60}$$

(c) Sketch the graph of $f(x)$, and sketch the linear approximation (i.e., the tangent line) at x_0 which you found in (a):



(d) Are the estimated values in part (b) overestimates or underestimates for the exact values for $f(x)$, i.e., is $L(x) > f(x)$ or is $L(x) < f(x)$? Give a brief explanation in terms of the graphs of $f(x)$ and $L(x)$ you sketched above.

Solution: We can see from the graph that the linear approximation (i.e., the tangent line) clearly sits above the curve—in other words, the y-values on the line are greater than the y-values on the curve (for any given value of x): $L(x) > f(x)$! Thus, the estimated values are overestimates.

3. (5 points) Read Problem 3 of the WebWork set, which describes a situation where a landscape architect wants to enclose a rectangular garden on one side by a brick wall and on the other three sides by metal fencing.
- (a) Using the variables and costs per foot for a brick wall and for metal fencing given in your WebWork, write down an expression for the total cost C in terms of x and y . Also draw a sketch of the garden, labeling it with the variables.

Solution:

Here is a sample WebWork exercise: "A landscape architect wished to enclose a rectangular garden on one side by a brick wall costing \$50/ft and on the other three sides by a metal fence costing \$20/ft. If the area of the garden is 122 square feet, find the dimensions of the garden that minimize the cost."

With these numbers, and if we let x = length of brick wall side and y = length of adjacent side, then the total cost is:

$$C(x, y) = 50x + 20y + 20y + 20x = 70x + 40y$$

(Obviously the brick wall will cost $50x$; the other 3 terms represent the cost of the 2 adjacent metal-fence sides of length y , and the metal-fence side opposite the brick side (so also of length x). Note that we have written the cost C here as a function of two variables, x and y .)

- (b) As asked in the WebWork exercise, solve for the dimensions of the garden that minimize the cost C .
Hint: use the "constraint" of the area of the garden given in the WebWork exercise to solve for y in terms of x , and substitute that into C in order to express the cost C as a function of only x . Then find the minimum of $C(x)$ by finding its critical point(s).

Solution: The constraint here is $xy = 122$, and so $y = \frac{122}{x}$. Substituting this for y in $C(x, y)$, we get cost as a function of x :

$$C(x) = 70x + 40 \left(\frac{122}{x} \right) = 70x + \frac{4880}{x}$$

Then:

$$C'(x) = 70 - \frac{488}{x^2} = \frac{70x^2 - 4880}{x^2}$$

So the critical points of $C(x)$ occur when $70x^2 - 4880 = 0$, i.e., for $x^2 = \frac{4880}{70}$. Hence, the maximal area occurs for $x = \sqrt{\frac{4880}{70}} \approx 8.35$ (feet)

To solve for the corresponding value of y , we use $y = \frac{122}{x}$, and so $y \approx \frac{122}{8.35} = 14.61$ (feet)

Entered	Answer Preview	Result
8.34951	$\sqrt{\frac{4880}{70}}$	correct
14.6116	$\frac{122}{\sqrt{\frac{4880}{70}}}$	correct

All of the answers above are correct.

A landscape architect wished to enclose a rectangular garden on one side by a brick wall costing \$50/ft and on the other three sides by a metal fence costing \$20/ft. If the area of the garden is 122 square feet, find the dimensions of the garden that minimize the cost.

Length of side with bricks $x = \sqrt{\frac{4880}{70}}$

Length of adjacent side $y = \frac{122}{\sqrt{\frac{4880}{70}}}$

4. (5 points) Find the derivative of $f(x) = 2x^2 - 7x + 1$ using the limit definition of the derivative, according to the following steps:

- (a) Write out and simplify $f(x + h)$:

$$\mathbf{Solution:} \quad f(x + h) = 2(x + h)^2 - 7(x + h) + 1 = 2(x^2 + 2xh + h^2) - 7x - 7h + 1 = 2x^2 + 4xh + 2h^2 - 7x - 7h + 1$$

- (b) Write out and simplify $f(x + h) - f(x)$:

$$\mathbf{Solution:} \quad f(x + h) - f(x) = (2x^2 + 4xh + 2h^2 - 7x - 7h + 1) - (2x^2 - 7x + 1) = 4xh + 2h^2 - 7h$$

- (c) Simplify the difference quotient $\frac{f(x + h) - f(x)}{h}$:

$$\mathbf{Solution:} \quad \frac{f(x + h) - f(x)}{h} = \frac{h(4x + 2h - 7)}{h} = 4x + 2h - 7$$

- (d) Finally, find $f'(x)$ by evaluating the limit:

$$\mathbf{Solution:} \quad f'(x) = \lim_{h \rightarrow 0} \frac{f(x + h) - f(x)}{h} = \lim_{h \rightarrow 0} (4x + 2h - 7) = 4x - 7$$

5. (5 points) Consider the function $f(x) = \frac{1}{3}x^3 - 3x^2 + 8x + 4$

(a) Find the first and second derivatives of $f(x)$:

Solution: $f'(x) = x^2 - 6x + 8$, $f''(x) = 2x - 6$

(b) Find the critical points of f , i.e., solve for the values x such that $f'(x) = 0$:

Solution: $f'(x) = x^2 - 6x + 8 = (x - 4)(x - 2) = 0 \implies x = 2, 4$

(c) For what values of x is $f'(x) > 0$ and for what values of x is $f'(x) < 0$? Show or explain how you solve for these intervals. What do these intervals represent in terms of the shape of the graph of $f(x)$?

Solution: Since $f'(x)$ is a quadratic with a positive leading term with roots at $x = 2$ and $x = 4$, we can conclude that $f'(x) > 0$ on $(-\infty, 2) \cup (4, \infty)$, meaning the graph is increasing on these intervals. On the other hand $f'(x) < 0$ on $(2, 4)$, and so the graph is decreasing on that interval.

It can help to draw a rough sketch of the graph, as I asked you to do on the in-class exercise.

(d) For what values of x is $f''(x) > 0$ and for what values of x is $f''(x) < 0$? Again, show or explain how you solve for these intervals, and explain what these intervals represent in terms of the shape of the graph of $f(x)$:

Solution: Since $f''(x) = 2x - 6$, which is a linear function with positive slope and x -intercept at $x = 3$, it is clear that $f''(x) > 0$ for all x in $(3, \infty)$ (and so the graph is concave up there) while $f''(x) < 0$ on $(-\infty, 3)$, where the graph is concave down. Hence, $x = 3$ is an inflection point.

(e) Sketch the graph of $y = f(x)$. Label the critical point(s) on the graph, and indicate whether each is a local maximum or a local minimum. Also label the y -intercept and any inflection points.

